

科技部補助專題研究計畫成果報告 期末報告

抗發炎機制在於Lumbrokinase心臟保護作用中扮演的角色(第 3年)

計畫類別：個別型計畫
計畫編號：MOST 103-2320-B-040-001-MY3
執行期間：105年08月01日至106年07月31日
執行單位：中山醫學大學醫學系藥理學科

計畫主持人：黃相碩
共同主持人：林素滿
計畫參與人員：碩士班研究生-兼任助理：洪岳慈
大專生-兼任助理：張家崧
大專生-兼任助理：周嘉宇
大專生-兼任助理：蔡仁維
大專生-兼任助理：姚懿庭
博士班研究生-兼任助理：王羿忻
博士班研究生-兼任助理：蘇倬慧

中華民國 106 年 08 月 03 日


中文摘要：由*Lumbricus rubellus*純化而來的抗血栓形成劑Lumbrokinase在臨床上已經被用於治療中風和心血管疾病。然而，Lumbrokinase的心臟保護作用與發炎反應的相關性仍然未知。在本研究中，我們在於心肌缺血再灌注損傷的大鼠。研究Lumbrokinase抑制發炎調節因子表現的訊息傳遞路徑。研究在麻醉大鼠的左主幹冠狀動脈進行1小時結紮和3小時再灌注傷害。動物分為不給予以及給予Lumbrokinase治療組，比較動物遭受心肌缺血再灌注損傷誘發的心律不齊和梗塞體積的嚴重程度。研究發現 Lumbrokinase抑制心肌缺血再灌注損傷誘發的心律不整並降低動物死亡率，並降低頸動脈血中的乳酸脫氫酶活性。Lumbrokinase經由TLR4訊息傳遞路徑抑制心肌缺血再灌注損傷增加的環氧合酶（COX）-2，誘導型一氧化氮合酶（iNOS）以及基質金屬蛋白酶（MMP）-9的蛋白質表現。此外，我們的研究結果證明，Lumbrokinase減少JNK，I κ B和NF- κ B的磷酸化。這些研究結果證明Lumbrokinase在於遭受心肌缺血再灌注損傷的大鼠是有效的心臟保護藥物，Lumbrokinase的心臟保護作用可能通過TLR4訊息路徑調節JNK和NF- κ B磷酸化路徑，進而減少心肌缺血再灌注損傷誘導COX-2，iNOS和MMP-9的表現相關。這些結果可以證明lumbrokinase是一個很有潛力的心臟保護劑，可以通過TLR4訊息路徑保護心臟對抗心肌缺血再灌注傷害。

中文關鍵詞：蚯蚓酵素Lumbrokinase；心臟保護；缺血；再灌注；TLR4

英文摘要：Lumbrokinase, a novel antithrombotic agent, purified from the earthworm *Lumbricus rubellus*, has been clinically used to treat stroke and cardiovascular diseases. However, inflammatory responses associated with the cardioprotective effect of lumbrokinase remain unknown. In this study, the signaling pathways involved in lumbrokinase-inhibited expressions of inflammation mediators were investigated in rats subjected to myocardial ischemia-reperfusion (I-R) injury. The left main coronary artery of anesthetized rats was subjected to 1 h occlusion and 3 h reperfusion. The animals were treated with/without lumbrokinase and the severities of I-R-induced arrhythmias and infarction were compared. Lumbrokinase inhibited I-R-induced arrhythmias and reduced mortality, as well as decreased the lactate dehydrogenase levels in carotid blood. Lumbrokinase also inhibited the enhancement of I-R induced expressions of cyclooxygenase (COX)-2, inducible nitric oxide synthase (iNOS), and matrix metalloproteinase (MMP)-9 through toll-like receptor 4 (TLR4) signaling pathway. Moreover, our results demonstrated that stimulation with lumbrokinase decreases the phosphorylation of JNK, IB, and NF- κ B. These findings suggested that lumbrokinase is a potent cardioprotective drug in rats with I-R injury. The cardioprotective effects of lumbrokinase may be correlated with its inhibitory effect on the I-R-induced expressions of COX-2, iNOS and MMP-9, mediated by TLR4 signaling

through JNK and NF- κ B pathways.

英文關鍵詞：Lumbrokinase; Cardioprotection; Ischemia; Reperfusion; TLR4



Lumbrokinase attenuates myocardial ischemia-reperfusion injury by inhibiting TLR4 signaling



Yi-Hsin Wang^a, Ke-Min Chen^b, Ping-Sung Chiu^b, Shih-Chan Lai^{b,c}, Hsing-Hui Su^d, Ming-Shiou Jan^{e,f,g}, Chia-Wei Lin^e, Dah-Yuu Lu^{h,i}, Yuan-Tsung Fu^{j,k}, Juan-Miaw Liao^{l,*}, Jinghua Tsai Chang^{a,m,**}, Shiang-Suo Huang^{a,n,o,***}

^a Institute of Medicine, Chung Shan Medical University, Taichung, Taiwan

^b Department of Parasitology, Chung Shan Medical University, Taichung, Taiwan

^c Department of Clinical Laboratory, Chung Shan Medical University Hospital, Taichung, Taiwan

^d Department and Institute of Pharmacology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^e Institute of Biochemistry, Microbiology and Immunology, Medical College of Chung Shan Medical University, Taichung, Taiwan

^f Division of Allergy, Immunology, and Rheumatology, Chung Shan Medical University Hospital, Taichung, Taiwan

^g Immunology Research Center, Chung Shan Medical University, Taichung, Taiwan

^h Department of Pharmacology, School of Medicine, China Medical University, Taichung, Taiwan

ⁱ Department of Photonics and Communication Engineering, Asia University, Taichung, Taiwan

^j School of Post-Baccalaureate Chinese Medicine, Tzu Chi University, Hualien, Taiwan

^k Department of Chinese Medicine, Taichung Tzu Chi Hospital, The Buddhist Tzu Chi Medical Foundation, Taichung, Taiwan

^l Department of Physiology, Chung Shan Medical University and Chung Shan Medical University Hospital, Taichung, Taiwan

^m Department of Medical Oncology and Chest Medicine, Chung Shan Medical University Hospital, Taichung, Taiwan

ⁿ Department of Pharmacology, Chung Shan Medical University, Taichung, Taiwan

^o Department of Pharmacy, Chung Shan Medical University Hospital, Taichung, Taiwan

ARTICLE INFO

Article history:

Received 1 December 2015

Received in revised form 17 July 2016

Accepted 4 August 2016

Available online 05 August 2016

Keywords:

Lumbrokinase
Cardioprotection
Ischemia
Reperfusion
TLR4

ABSTRACT

Lumbrokinase, a novel antithrombotic agent, purified from the earthworm *Lumbricus rubellus*, has been clinically used to treat stroke and cardiovascular diseases. However, inflammatory responses associated with the cardioprotective effect of lumbrokinase remain unknown. In this study, the signaling pathways involved in lumbrokinase-inhibited expressions of inflammation mediators were investigated in rats subjected to myocardial ischemia-reperfusion (I-R) injury. The left main coronary artery of anesthetized rats was subjected to 1 h occlusion and 3 h reperfusion. The animals were treated with/without lumbrokinase and the severities of I-R-induced arrhythmias and infarction were compared. Lumbrokinase inhibited I-R-induced arrhythmias and reduced mortality, as well as decreased the lactate dehydrogenase levels in carotid blood. Lumbrokinase also inhibited the enhancement of I-R induced expressions of cyclooxygenase (COX)-2, inducible nitric oxide synthase (iNOS), and matrix metalloproteinase (MMP)-9 through toll-like receptor 4 (TLR4) signaling pathway. Moreover, our results demonstrated that stimulation with lumbrokinase decreases the phosphorylation of JNK, IκB, and NF-κB. These findings suggested that lumbrokinase is a potent cardioprotective drug in rats with I-R injury. The cardioprotective effects of lumbrokinase may be correlated with its inhibitory effect on the I-R-induced expressions of COX-2, iNOS and MMP-9, mediated by TLR4 signaling through JNK and NF-κB pathways.

© 2016 Published by Elsevier Ltd.

1. Introduction

Coronary artery and associated ischemic heart diseases are prevalent worldwide. Myocardial ischemia-reperfusion (I-R) injury, which is one

of the world's fastest growing conditions, is a major contributor to the morbidity and mortality associated with coronary artery diseases [1,2]. Myocardial ischemia is a heart condition resulting from lack of coronary blood supply caused by thrombosis or acute alterations of coronary

Abbreviations: I-R, ischemia-reperfusion; COX-2, cyclooxygenase-2; iNOS, inducible nitric oxide synthase; MMP-9, matrix metalloproteinase-9; TLR4, toll-like receptor 4; e-PA, extrinsic plasminogen activator; t-PA, tissue plasminogen activator; HR, heart rate; BP, arterial blood pressure; VPC, ventricular premature complexes; VT, ventricular tachycardia; VF, ventricular fibrillation; LDH, lactate dehydrogenase; MAPKs, mitogen-activated protein kinases.

* Correspondence to: J.-M. Liao, Department of Physiology, Chung Shan Medical University, Taichung, No. 110, Sec. 1, Jianguo N. Rd., Taichung City 402, Taiwan.

** Correspondence to: J. T. Chang, Institute of Medicine, Chung Shan Medical University, Taichung, No. 110, Sec. 1, Jianguo N. Rd., Taichung City 402, Taiwan.

*** Correspondence to: S.-S. Huang, Department of Pharmacology and Institute of Medicine, Chung Shan Medical University, and Department of Pharmacy, Chung Shan Medical University Hospital, Taichung, No. 110, Sec. 1, Jianguo N. Rd., Taichung City 402, Taiwan.

E-mail addresses: jmliao@csmu.edu.tw (J.-M. Liao), jinghuat@csmu.edu.tw (J.T. Chang), sshuang@csmu.edu.tw (S.-S. Huang).

atherosclerotic plaques [1]. During myocardial ischemia, a characteristic pattern of metabolic and ultrastructural changes leads to irreversible injury. Early restoration of blood flow to the ischemic myocardium is a common treatment strategy to limit myocardial infarct size and to reduce mortality [3]. However, the return of blood flow can cause additional cardiac damage and complications that can paradoxically increase infarct size, a condition referred to as myocardial I-R injury [4,5].

Myocardial I-R injury is a pathological condition characterized by an initial restriction of blood supply to the myocardium, subsequent restoration of perfusion, and concomitant reoxygenation. Over the past two decades, studies on myocardial I-R injury have focused on the role of molecular, rather than cellular, targets such as complement receptors [6,7], toll-like receptors (TLRs) [8] and proinflammatory cytokines [7]. Myocardial I-R injury is characterized by a rapid increase in cytokines and chemokines and an influx of leukocytes into the vulnerable region bordering the infarcted site. This inflammatory response may not only lead to cardiomyocyte apoptosis, but also compromise myocardial function. Therefore, limiting I-R-induced myocardial inflammation may lower mortality and reduce myocardial I-R-induced ventricular arrhythmia and myocardial infarction [9].

Lumbrokinase, an extract of *Lumbricus rubellus*, was identified more than two decades ago [10]. Lumbrokinase is a group of bioactive proteolytic enzymes with molecular weights ranging from 25 kDa to 32 kDa [11]. Lumbrokinase includes plasminogen activator and plasmin [12]. The extrinsic plasminogen activator (e-PA) in lumbrokinase is similar to the tissue plasminogen activator (t-PA) found in other species. However, lumbrokinase dissolves fibrin clots by converting plasminogen to plasmin, while exhibiting relatively good heat stability and broad optimal pH range [13]. Lumbrokinase enzymes demonstrate thrombolytic activity only in the presence of fibrin. Therefore, lumbrokinase does not cause hemorrhage, as opposed to streptokinase or urokinase, due to hyperfibrinolysis during treatment [12]. Lumbrokinase has been clinically applied to the treatment of strokes and cardiovascular diseases [14]. In China, lumbrokinase has been marketed for >10 years mainly for the treatment of acute-phase ischemic stroke [14], prevention of secondary ischemic stroke [15], and improvement of myocardial perfusion in patients with stable angina [16]. However, few studies have reported the pharmacological mechanisms of the effects of lumbrokinase on myocardial I-R injury in rats.

Evidence has shown that TLRs play a central role in I-R injury [8,17,18]. The results of the present study demonstrated that lumbrokinase treatment significantly attenuates the myocardial I-R-induced expression of the TLR4 protein when compared with vehicle-treated group. It is well known that TLR4 signaling plays a dominant role in myocardial I-R injury [8]. Mice lacking TLR4 sustain smaller infarct size and exhibit less inflammation when compared with wild-type mice after myocardial I-R injury [18]. Therefore, modulation of TLR4 signaling may be a potential therapeutic strategy in myocardial I-R injury [19]. The present study aimed to explore the possible anti-inflammatory mechanisms (with specific focus on the TLR4 signaling pathway) of lumbrokinase for ameliorating myocardial I-R injury in rats.

2. Methods

Study protocols conformed with those published in the Guide for the Care and Use of Laboratory Animals by the National Research Council of the National Academies (NIH publication, revised 2011). Every effort was made to minimize animal suffering and to reduce the number of sacrificed animals.

2.1. Animals

Six-week old male Sprague-Dawley rats (National Laboratory Animal Center) weighing 250–300 g were housed in the Animal Center of Chung Shan Medical University at an ambient temperature of

25 ± 1 °C, under a 12 h light–dark cycle. The animals were fed normal chow and given water ad libitum. All surgical procedures for myocardial I-R were reviewed and approved by the Institutional Animal Care and Use Committee of Chung Shan Medical University.

2.2. Experimental protocol

Myocardial I-R injury was induced by temporary occlusion of the left main coronary artery as previously described [20]. Briefly, the rats were anesthetized with a single intraperitoneal injection of thiobutobarbital sodium salt (100 mg/kg) and urethane (400 mg/kg) and then placed on a heated, small animal operating table. Rectal temperature was maintained at 38 ± 0.5 °C. After tracheotomy, the intubated animals were ventilated with room air through a respirator for small rodents (Model 131, NEMI, USA) with stroke volume of 15 ml/kg body weight and rate of 60 strokes/min to maintain normal P_{O₂}, P_{CO₂} and pH parameters (blood gas analyzer, GEM-5300 IL, CO, USA). The jugular vein was cannulated to administer drugs and Evans blue at the end of the experiment. Lumbrokinase (1 or 10 µg/kg, Canada RNA Biochemical Inc., Richmond, BC, Canada) or vehicle (normal saline) was infused via the jugular vein 15 min before coronary artery occlusion. The rats injected with vehicle were used as controls. The vehicle had no effects on myocardial I-R-induced arrhythmia or infarction. The animals were randomly allocated to drug treatment or vehicle administration groups. Polyethylene catheters (PE-50) were inserted into the common carotid artery with a Statham P23 XL transducer to continuously monitor heart rate (HR) and arterial blood pressure (BP), as well as adequacy of the anesthesia. Data were acquired with a Gould RS-3400 physiological recorder (Gould, Cleveland, OH, USA). Standard lead-1 ECG was recorded via silver electrodes attached to the extremities.

The heart was exposed via left thoracotomy and the fourth and fifth ribs were sectioned approximately 2 mm to the left of the sternum. The heart was rapidly externalized and inverted. Then, a 6/0 silk ligature was placed around the left main coronary artery. The heart was repositioned in the chest and the animal was allowed to recover for 15 min. Animals that developed arrhythmia or sustained a decrease in mean BP to <70 mmHg during the procedure were not included in the study. A small plastic snare from a Portex P-270 cannula was threaded through the ligature and placed in contact with the heart. The left coronary artery was then occluded by tightening the ligature and reperfusion was achieved by releasing the tension applied to the ligature (operated groups). Successful ligation of the coronary artery was validated by decreased arterial pressure and ECG changes (increase in R wave and ST segment elevation), which are indicators of ischemia. Sham-operated animals underwent the same surgical procedure, except that the silk ligature that was placed around the left coronary artery was not tied.

Changes in HR, BP, and ECG were simultaneously recorded using a personal computer with waveform analysis software (AcqKnowledge, Biopac System, Goleta, CA, USA) before and during the ischemia and reperfusion periods. To evaluate antiarrhythmic effects of lumbrokinase during myocardial I-R injury, we occluded the coronary artery for 1 h and then subjected it to reperfusion for 3 h. The ventricular ectopic activity was evaluated according to the diagnostic criteria advocated by the Lambeth Conventions [21]. Incidence and duration of ventricular tachyarrhythmias, including ventricular tachycardia (VT) and ventricular fibrillation (VF), were determined in the surviving rats and in the rats that eventually died. In rats with irreversible VF, the duration of VF was recorded until mean BP was <15 mmHg.

2.3. Determination of infarct size and collection of myocardium samples

Evaluation of the infarct zone and collection of myocardium samples for further analyses were carried out in rats that survived 1 h of ischemia and 3 h of reperfusion. To reduce the number of animals used in this study, an effective dose of 10 µg/kg was applied to investigate whether

lumbrokinase ameliorates myocardial I-R injury and to uncover the underlying mechanism involved in the cardioprotective effect of lumbrokinase. The sizes of the occluded and infarct zones in the hearts were determined using a previously described procedure [22]. At the end of the experiment, the coronary artery was reoccluded, and 2.0 ml of 3% Evans blue solution were intravenously injected to stain non-ischemic myocardium and determine the area at risk. The heart was then excised and the atria were removed. Ventricular tissues were sliced into 1 mm sections and incubated in tetrazolium dye [2% 2,3,5-triphenyltetrazolium chloride (Sigma, USA) in normal saline] at 37 °C for 40 min in darkness. The sections were placed in a solution of 10% formaldehyde in saline for 1 day before excision of infarcted (white) tissues. The weight of the infarcted tissues is expressed as a percentage of the total ventricle or the area at risk.

2.4. Determination of myocardial damage

Myocardial cellular damage was estimated by measuring lactate dehydrogenase (LDH) leakage into plasma. Arterial blood was collected from the carotid catheter at the end of myocardial I-R injury for LDH measurements. LDH activity was determined using a commercially available assay kit (Sigma, St Louis, MO, USA).

2.5. Protein extraction and western blot analysis

Heart tissues were homogenized with a tissue protein extraction reagent (Thermo Scientific, USA) containing protease inhibitor cocktail (Sigma, USA). The tissue homogenates were centrifuged at 12,000 g for 10 min at 4 °C. Protein concentration was determined with protein assay kits (BioRad, USA) using bovine serum albumin as the standard. The samples were mixed with an equal volume of loading buffer [62.5 mM Tris-HCl, pH 6.8, 10% (v/v) glycerol, 2% SDS, 5% (v/v) 2-mercaptoethanol, and 0.05% (w/v) bromophenol blue] and heated at 95 °C for 5 min.

The protein samples were subjected to SDS-PAGE and electrophoretically transferred onto PVDF protein sequencing membranes for 90 min. The membrane was blocked with 5% non-fat milk in PBS-buffered saline with 0.1% (v/v) Tween-20 (PBST) at room temperature for 1 h. The membrane was then washed and blotted with the antibodies of matrix metalloproteinase (MMP)-9, MMP-2 (R&D Systems, USA), COX-2 (Cayman Chemical, USA), TLR4, TLR2, c-fos (Santa Cruz, USA), interleukin-1 receptor-associated kinase (IRAK)4, phospho-IRAK4, phospho-JNK, JNK, phospho-NF- κ B, NF- κ B, phospho-I κ B- α , I κ B- α , iNOS, eNOS, nNOS, phospho-ERK 1/2, ERK 1/2, phospho-p38, and p38 (Cell Signaling, USA), MPO (Spring Bioscience, USA). β -actin (Abcam, UK) served as internal loading control. The membrane was incubated with HRP-conjugated secondary antibody (Jackson ImmunoResearch Laboratories, USA) prior to chemiluminescence detection (Pierce, USA).

2.6. Gelatin zymography

Myocardial I-R-induced proteinase in rat hearts was analyzed by gelatin zymography as previously described [23] with slight modification. Briefly, the heart homogenates were loaded onto SDS-PAGE (7.5%) polyacrylamide gel, which was co-polymerized with 0.1% (w/v) gelatin (Sigma, USA). The stacking gels were comprised of 4% (w/v) polyacrylamide without gelatin substrate. Electrophoresis was performed in running buffer (25 mM Tris, 250 mM glycine, 1% SDS) at room temperature. The gel was washed twice in double-distilled water containing 2.5% Triton X-100 for 30 min each time, then incubated in reaction buffer [50 mM Tris-HCl, pH 7.5, 200 mM NaCl, 10 mM CaCl₂, 0.02% (w/v) Brij@-35, 0.01% (w/v) NaN₃] at 37 °C for 18 h. Subsequently, the gel was stained with 0.25% (w/v) Coomassie brilliant blue R-250 (Sigma, USA) for 1 h and destained with 15% (v/v) methanol and 7.5% (v/v) acetic acid. Gelatinase activity was detected as unstained bands on a blue background. Quantitative analysis was performed with a

computer-assisted imaging densitometer system (UN-SCAN-IT™ gel version 5.1, Silk Scientific, UT).

2.7. Immunohistochemical staining

Rat hearts were fixed separately in 10% neutral-buffered formalin for 24 h. They were then dehydrated in a graded ethanol series (50%, 75%, 95% and 100%), cleared in xylene, and embedded in paraffin at 55 °C for 24 h. For the immunohistochemical analysis, relatively thick (5 μ m) serial sections of wax-embedded heart were cut, mounted on glass slides, dewaxed, and then rehydrated in the same manner as for histological analysis. Subsequently, the sections were treated with 3% H₂O₂ in methanol for 10 min to inactivate any endogenous peroxidase and washed thrice with PBS for 5 min each time. Afterward, the sections were blocked with 3% BSA at room temperature for 1 h and then incubated at 37 °C for 1 h with a 1:50 dilution of the rabbit anti-human MPO polyclonal antibody and a 1:100 dilution of the rabbit anti-human TLR4 polyclonal antibody in 1% BSA. Following another three washes in PBS, the sections were incubated at 37 °C for 1 h with a 1:100 dilution of the HRP-conjugated rabbit anti-goat IgG in 1% BSA and then washed three more times in PBS. Finally, the sections were incubated at room temperature for 3 min with 3,3'-diaminobenzidine (0.3 mg/ml) in 100 mM Tris (pH 7.5) containing 0.3 μ l H₂O₂/ml. After washing thrice in PBS, the sections were mounted in 50% glycerol in PBS and examined under light microscope.

2.8. Flow cytometry analysis

The peripheral blood mononuclear cells (PBMCs) were washed twice with 3 ml of FACScan medium (RPMI-1640 medium with 2% FBS and 0.1% NaN₃). An aliquot of 1×10^5 cells was used for each FACScan tube (FALCON 352052). After double labeling with either a mixture of anti-TLR4-PE (Novus Biologicals, USA) and anti-CD3 FITC (BD Biosciences, USA) or a mixture of anti-TLR4-PE and anti-CD19-FITC (Abcam, UK) for 45 min on ice in the dark, cells were washed twice and resuspended in FACScan medium. The fluorescence intensity was detected with flow cytometer (FACScalibur; Becton Dickinson, USA) and analyzed with Flowing Software (2.5.1). The expressions of TLR4 in T lymphocytes and B lymphocytes are presented as the values of geometric mean fluorescence intensity (gMFI).

2.9. Blood-clot lysis activity assay

An in vitro blood-clot lysis activity assay was performed as described by Prasad et al. Fresh rat blood was drawn from the carotid artery. Then, 600 μ l of blood were transferred to pre-weighed Eppendorf tubes and incubated at 37 °C for 45 min. Serum was completely removed after clot formation, and the tubes containing the clots were weighed again. Clot weight was determined by subtracting the weight of each tube (i.e., clot weight = weight of clot-containing tube – weight of empty tube). Then, 200 μ l of lumbrokinase at varying concentrations (10, 100, 96,000 units), dissolved in PBS, were added to the tubes containing clots. The tubes were then incubated at 37 °C for 90 min and observed for clot lysis. Lysed fluid was completely absorbed from each tube with filter paper and the tubes were then re-weighed. Weight differences of each tube before and after incubation were calculated for treated and control samples.

2.10. Statistical analysis

Data are expressed as mean \pm standard error of mean (SEM). Statistical analyses of differences were carried out using one-way analysis of variance (ANOVA) for combined data followed by Bonferroni's tests. The differences in the percentage incidences of VT, VF, and mortality were compared using χ^2 test. Spearman rank correlation was applied to test the correlations between MMP-9 activity and MPO expression

in rats subjected to myocardial I-R injury. $p < 0.05$ was considered statistically significant.

3. Results

3.1. Hemodynamics

There were no statistically significant differences in pre-ischemic hemodynamic parameters among the groups of rats. The injection of lumbrokinase into the jugular vein did not affect the mean BP or HR of rats. Moreover, hemodynamic parameters did not significantly differ between the vehicle- and lumbrokinase-treated rats during myocardial I-R injury (Fig. 1).

3.2. Myocardial I-R-induced mortality and rhythm disturbances

The effects of lumbrokinase on myocardial I-R-induced mortality and arrhythmia in rats are shown in Table 1. In the vehicle-treated group, the I-R-induced mortality was 53.3%. However, administration of 10 $\mu\text{g}/\text{kg}$ lumbrokinase significantly reduced the I-R-induced mortality to 10%.

Jugular vein injection of the vehicle or lumbrokinase did not elicit arrhythmia in rats. The number of ventricular premature complexes (VPC) was 53.8 ± 28.4 , and the durations of VT and VF were 19.0 ± 4.8 and 82.9 ± 14.9 s, respectively, in the vehicle-treated group during the I-R period. However, administration of 10 $\mu\text{g}/\text{kg}$ lumbrokinase significantly decreased the number of VPC to 16.7 ± 7.0 and the durations of VT and VF to 3.0 ± 0.8 and 2.3 ± 1.3 s, respectively.




Fig. 1. (A) Mean BP and (B) heart rate (C) ECG were recorded during myocardial I-R injury. Each experimental group contains ten rats. Results are expressed as mean \pm SEM. No significant differences were recorded between the vehicle and two doses of lumbrokinase (LK-1: 1 $\mu\text{g}/\text{kg}$; LK-10: 10 $\mu\text{g}/\text{kg}$) treated rats. L indicates coronary ligation. R indicated reperfusion.

3.3. Myocardial damage and infarct size

There were no significant differences in the size of the area at risk between the vehicle- and lumbrokinase-treated rats, indicating that similar amounts of myocardial tissues were at risk from occlusion of the left coronary artery in each group. In the vehicle-treated group, the infarct size was $47.4 \pm 1.8\%$ of the area at risk. However, administration of $10 \mu\text{g}/\text{kg}$ lumbrokinase significantly reduced the infarct size to $19.6 \pm 1.4\%$ of the area at risk. Furthermore, in comparison with vehicle-treated rats, administration of $10 \mu\text{g}/\text{kg}$ lumbrokinase significantly decreased LDH activity, a marker of cell damage, after myocardial I-R injury. The resulting LDH levels were consistent with infarct size data (Table 2).

3.4. Effects of lumbrokinase on the protein expressions of TLR2 and TLR4 in myocardial tissues

TLRs play a central role in I-R injury [8,17,18] and may be involved in the cardioprotective effects of lumbrokinase (Fig. 2). As shown in Fig. 2A and B, TLR2 expression did not significantly change due to vehicle or $10 \mu\text{g}/\text{kg}$ lumbrokinase treatment in rats subjected to myocardial I-R injury. However, the protein expression of TLR4 significantly increased in the vehicle-treated group when compared with the sham-operated group. In contrast, lumbrokinase treatment significantly attenuated the myocardial I-R-induced expression of the TLR4 protein in comparison with the vehicle-treated group.

Immunohistochemical staining for TLR4 was performed on the tissues acquired from the area at risk. Lumbrokinase treatment reduced TLR4 staining in the myocardium obtained from rats subjected to myocardial I-R (Fig. 2C–G) when compared with the vehicle group. This is consistent with our finding that pretreatment with lumbrokinase decreases the protein expression of TLR4 in myocardial tissues after I-R injury. As TLRs are also expressed in immune cells [24], we conducted experiments using flow cytometry to determine TLR4 expression in lymphocytes following myocardial I-R injury. Supplementary Fig. 1 shows similar levels of expressions of TLR4 in T lymphocytes between lumbrokinase-treated and vehicle-treated groups. The expression of TLR4 was lower in B lymphocytes in the lumbrokinase-treated group when compared with the vehicle-treated group, however the difference did not reach statistical significance. Fallach et al. reported that cardiac function of TLR4-knockout mice and chimeric mice expressing TLR4 in the immunohematopoietic system, but not in the heart, is resistant to LPS and reduced cardiac depression following MI [24]. In addition, Avlas et al. observed that cardiac function is significantly less depressed following coronary artery ligation in chimeric mice deficient in TLR4 in the heart, but not in the immunohematopoietic system, similar to TLR4-knockout mice [25]. These findings suggest that TLR4 expressed by the cardiomyocytes themselves plays a key role in acute myocardial dysfunction caused by septic shock and myocardial ischemia [24,25]. Moreover, TLR4/NF- κ B signal pathway is known to be of importance in I-R injury and to be markedly upregulated in ischemic myocardium [26].

Table 1

Effect of lumbrokinase on arrhythmias induced by myocardial I/R injury in anesthetized rats.

	n	VPC		VT		VF		Mortality (%)
		Incidence (%)	Count (number)	Incidence (%)	Duration (sec)	Incidence (%)	Duration (sec)	
Sham		–	–	–	–	–	–	–
Vehicle	5	–	–	–	–	–	–	–
LK-10	5	–	–	–	–	–	–	–
Operated (ligated)								
Vehicle	15	80	53.8 ± 28.4	67	19.0 ± 4.8	47	82.9 ± 14.9	53.3
LK-1	10	50	34.6 ± 9.5	40	$6.2 \pm 1.1^*$	10*	$4.7 \pm 2.1^*$	20
LK-10	10	30*	$16.7 \pm 7.0^*$	20*	$3.0 \pm 0.8^*$	10*	$2.3 \pm 1.3^*$	10*

VPC, ventricular premature complexes; VT, ventricular tachycardia; VF, ventricular fibrillation.

Vehicle is normal saline. n, number of experiments; values for duration of VT and VF are shown as the mean \pm SEM.

* Statistical difference at the level of $p < 0.05$ as compared with vehicle.

Table 2

Weight and size of area at risk ($n = 6$).

	Vehicle	LK-10
LV weight (g)	0.94 ± 0.05	0.87 ± 0.06
Area at risk (g)	0.46 ± 0.03	0.43 ± 0.03
Area at risk/LV (%)	49.2 ± 0.6	49.4 ± 1.4
Infarct size (g)	0.21 ± 0.04	$0.08 \pm 0.01^*$
Infarct size/LV (%)	23.4 ± 6.2	$9.7 \pm 0.8^*$
Risk zone infarcted (%)	47.4 ± 1.8	$19.6 \pm 1.4^*$
Lactate dehydrogenase (U/L)	6197 ± 1833	$1856 \pm 53^*$

LV, left ventricle.

Data are shown as the mean \pm SEM.

* Statistical difference at the level of $p < 0.05$ as compared with vehicle.

As shown in Fig. 2E and F, intense TLR4 staining was observed in the cardiomyocytes of vehicle-treated group. However, TLR4 positive staining was rarely observed in the cardiomyocytes of lumbrokinase-treated group. These results are consistent with the findings of previous reports.

3.5. Effects of lumbrokinase on TLR4 signaling pathway

Fig. 3 shows the effects of lumbrokinase on TLR4 signaling pathway in rats subjected to myocardial I-R injury. In parallel with the TLR4 expression, IRAK4 protein expression and phosphorylation level were also examined. The expression of the IRAK4 protein was not altered by vehicle or lumbrokinase treatment in rats subjected to myocardial I-R injury. However, the phosphorylated IRAK4 (p-IRAK4) protein level significantly increased in the vehicle-treated group. This increase was alleviated by treatment with lumbrokinase (Fig. 3A and B).

We studied the signaling pathways involved in the cardioprotective effect of lumbrokinase. The protein expressions of p38, ERK 1/2 and JNK and their phosphorylation levels and c-fos were determined. As shown in Fig. 3C–G, treatment with vehicle or lumbrokinase did not affect the expressions of p38, phosphorylated p38 (p-p38), ERK 1/2, phosphorylated ERK 1/2 (p-ERK 1/2), or JNK. However, phosphorylated JNK (p-JNK) and c-fos expressions were higher in the vehicle-treated group. Lumbrokinase treatment diminished the increases in p-JNK and c-fos expressions. Additionally, the protein expressions of phosphorylated I κ B (p-I κ B), NF- κ B, and phosphorylated NF- κ B (p-NF- κ B) significantly increased in the vehicle-treated group when compared with the sham-operated group. In contrast, lumbrokinase treatment significantly attenuated the myocardial I-R-induced increases in expressions of p-I κ B, NF- κ B, and p-NF- κ B when compared with the vehicle-treated group (Fig. 3H–J).

3.6. Effects of lumbrokinase on the protein expressions of NOSs, COX-2, and MMPs activity in myocardial tissues

The effects of lumbrokinase on the protein expressions of iNOS, eNOS, and nNOS in rats subjected to myocardial I-R injury are shown




Fig. 2. Effects of lumbrokinase on TLRs protein expression in rats subjected to myocardial I-R injury. Rats were treated with vehicle or 10 $\mu\text{g}/\text{kg}$ lumbrokinase (LK-10). (A) Representative images of the Western blot results TLR2 and TLR4 expression in heart tissue samples from rats after myocardial I-R injury. (B) Quantitative densitometric analysis of TLR2 and TLR4 protein expression with β -actin as an internal standard. Data were normalized with sham and presented as percentage rates. Each value represents the mean of three individual experiments. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with sham. # $p < 0.05$ compared with vehicle. (C–F) Representative areas of TLR4 immunohistochemical (IHC) staining slides of rats subjected to myocardial I-R injury. Panels C and D original magnification, panels E and F are higher magnification of small box in panels C and D. The dark brown color indicates expression of TLR4. (G) Quantitative densitometric analysis of TLR4 staining slides of rats subjected to myocardial I-R injury. Data were normalized with vehicle and presented as percentage rates. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with vehicle.

in Fig. 4A. The protein expressions of eNOS and nNOS were not altered by vehicle or lumbrokinase treatment in rats subjected to myocardial I-R injury. However, the protein expression of iNOS in the vehicle-treated group significantly increased when compared with the sham-operated group. In contrast, lumbrokinase treatment significantly attenuated the myocardial I-R-induced increase in expression of iNOS protein when compared with the vehicle-treated group. In addition, the expression of COX-2 significantly increased in the vehicle-treated group when compared with the sham-operated group. However, marked increase in COX-2 expression in rats with myocardial I-R was prevented by treatment with lumbrokinase (Fig. 4B).

Zymography was performed to measure the changes in the activities of MMP-2 and MMP-9, which are the characteristic MMPs of the myocardium [27]. The effects of lumbrokinase on the activities of MMP-2 and MMP-9 in rats subjected to myocardial I-R injury are shown in Fig. 4C. MMP-2 activity was not altered following vehicle or lumbrokinase treatment. However, MMP-9 activity significantly increased in the vehicle-treated group when compared with the sham-operated group. Administration of lumbrokinase significantly decreased the activity of MMP-9 in rats after myocardial I-R injury.

3.7. Effects of lumbrokinase on myocardial inflammation

Myocardial TLR4 and MMP-9 are critical to the myocardial expressions of chemoattractant proteins that mediate neutrophil infiltration [28]. We further examined whether smaller infarction in the

lumbrokinase-treated group corresponds to less myocardial inflammation as demonstrated by neutrophil infiltration. The protein expression of MPO, a marker enzyme of neutrophils, significantly increased in the vehicle-treated group when compared with the sham-operated group. In contrast, lumbrokinase treatment significantly attenuated the myocardial I-R-induced increase in expression of the MPO protein when compared with the vehicle-treated group (Fig. 5A and B). Immunohistochemical staining for MPO was performed on the tissues obtained from the area at risk. Hearts from the vehicle-treated group exhibited more neutrophil infiltration (arrow). Administration of lumbrokinase significantly reduced the I-R induced neutrophil infiltration into heart tissues when compared with the vehicle group (Fig. 5C–G). Therefore, we suggest that lumbrokinase decreases TLR4 protein expression and MMP-9 activity, which in turn reduces inflammation of myocardial tissues after I-R injury.

4. Discussion & conclusions

The cardioprotective effect of lumbrokinase in rats subjected to myocardial I-R injury was examined. Administration of 10 $\mu\text{g}/\text{kg}$ lumbrokinase significantly suppressed the incidence and number of VPCs, as well as the incidences and durations of VT and VF. This treatment also reduced mortality and size of cardiac infarct zone in rats subjected to 1 h of coronary artery occlusion and 3 h of reperfusion. These results are consistent with our finding that pretreatment with lumbrokinase leads to decreased carotid blood LDH levels during I-R





Fig. 3. Effects of lumbrokinase on TLR4 signaling pathway in rats subjected to myocardial I-R injury. Rats were treated with vehicle or 10 $\mu\text{g}/\text{kg}$ lumbrokinase (LK-10). (A) Representative images of the Western blot results. (B) Quantitative densitometric analysis of IRAK4 and p-IRAK4. (C) Representative images of the Western blot results. Quantitative densitometric analysis of (D) p38 and p-p38 (E) ERK1/2 and p-ERK1/2 (F) JNK and p-JNK (G) c-fos (H) Representative images of the Western blot results. Quantitative densitometric analysis of (I) I κ B and p-I κ B (J) NF- κ B and p-NF- κ B protein expression in heart tissue samples from rats after myocardial I-R injury. Data were normalized with sham and presented as percentage rates. Each value represents the mean of three individual experiments. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with sham. # $p < 0.05$ compared with vehicle.

injury. LDH level is an indicator of cellular damage. In addition, we demonstrated that the cardioprotective effect of lumbrokinase in rats after myocardial I-R injury via attenuated TLR4 regulates signaling pathway. Lumbrokinase exhibited inhibitory effects on myocardial I-R-activated TLR4 and activation of the corresponding downstream signaling molecules such as IRAK4, JNK, c-fos, I κ B, and NF- κ B. Subsequently, myocardial I-R-induced upregulation of MMP-9 activation and iNOS and COX-2 protein expressions were reduced, thus inhibiting myocardial I-R injury-induced inflammation.

TLR4 signaling plays a dominant role in myocardial I-R injury. TLR4 expression and infiltrating leukocytes within the myocardium significantly increase after I-R injury [8]. Accumulating evidence supports the notion that activation of the TLR4 pathway directly impairs the contractility of isolated cardiomyocytes [24,25,26,29]. Furthermore, mice lacking TLR4 sustain smaller infarct size and exhibit less inflammation when compared with wild-type mice after myocardial I-R injury [18]. Therefore, modulation of TLR4 signaling may be a potential therapeutic target in myocardial I-R injury [19]. Myeloid differentiation factor 88



have strongly suggested that TLR4 activation confers potent cardiac protection against I-R injury via an MyD88-dependent, Trif-independent signaling mechanism [30,31]. Recent studies have reported that TLR4-mediated NF- κ B signaling contributes to myocardial I-R injury while TLR4-mediated MyD88-dependent signaling pathway is essential for NF- κ B activation [32]. Activated MyD88 recruits IRAK4 and IRAK1, leading to the phosphorylated degradation of I κ B and NF- κ B activation [33]. The present study demonstrated that myocardial I-R injury activates IRAK4 through TLR4 signaling. Importantly, lumbrokinase inhibited the myocardial I-R-induced increase in TLR4 expression and markedly reduced IRAK4 activation. After TLR4 activation, the downstream signaling molecules, including transcription factor NF- κ B and mitogen-activated protein kinases (MAPKs), can be activated [34]. In this study, lumbrokinase effectively inhibited myocardial I-R-induced phosphorylation/activation of JNK and c-fos expressions. The JNK/c-fos pathway has been implicated in the signal transduction pathways responsible for the regulation of inflammatory response following myocardial I-R injury [35,36]. Although studies have shown that ERK1/2 and p38 MAPK are also involved in the production of inflammatory mediators, lumbrokinase did not reduce their production or phosphorylation in rats subjected to myocardial I-R injury in the present study. These observations indicated that JNK/c-fos, but not ERK1/2 and p38, MAPK-dependent pathways, are involved in the cardioprotective effect of lumbrokinase in rats after myocardial I-R injury.

The activation of transcription factor NF- κ B and MAPKs results in the induction of COX-2 and iNOS expressions. COX-2 and iNOS are co-existing, key inflammatory proteins that play fundamental roles under similar pathophysiological conditions [37]. In this study, lumbrokinase reduced the protein expressions of COX-2 and iNOS in rats after myocardial I-R injury. Lumbrokinase inhibited myocardial I-R-induced activation of the NF- κ B signaling cascade and phosphorylation of JNK, rather than p38 MAPK and ERK1/2. This suggested that the NF- κ B and JNK pathways modulate the suppression of the myocardial I-R-induced protein expressions of COX-2 and iNOS by lumbrokinase.

In addition, we examined inflammatory cell infiltration into the ischemic zone of the I-R heart tissues. The results of the present study demonstrated that lumbrokinase reduces neutrophil infiltration into the injured myocardial tissues after I-R. Accumulating evidence has demonstrated that infiltrating neutrophils are the major source of MMP-9 and may localize MMP-9 activation in the I-R injured myocardium [38,39]. A significant positive correlation was found between the relative intensity of MMP-9 and number of MPO positive cells in myocardium ($r = 0.7921$, $p < 0.05$), as shown in Supplementary Fig. 3. In addition, previous studies have demonstrated increases in the expression and activity of myocardial MMP-9 in experimental myocardial I-R injury. Thus, MMP-9 may serve as a target for prevention or treatment of acute myocardial I-R injury [38,39,40]. The results also showed that lumbrokinase reduces inflammation during myocardial I-R, thus limiting the infiltration of neutrophils into the injured tissues. Moreover, MMP-9 activity significantly decreased in the lumbrokinase-treated group when compared with the vehicle-treated group after myocardial I-R injury.

Lumbrokinase has been used in oral dosage to prevent and treat clotting, such as that related to myocardial infarction and cerebral thrombus [41]. Oral lumbrokinase (300,000 units) administered three

Fig. 4. Effects of lumbrokinase on the protein expression of NOSs and COX-2 and MMP-9 activity in rats subjected to myocardial I-R injury. Western blot analysis for (A) NOSs (B) COX-2 expression, and gelatin zymography analysis for (C) MMP-2 and MMP-9 activity in heart tissue samples from rats after myocardial I-R injury. Rats were treated with vehicle or 10 μ g/kg lumbrokinase (LK-10). Data were normalized with sham and presented as percentage rates. Each value represents the mean of three individual experiments. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with sham. # $p < 0.05$ compared with vehicle.

(MyD88) and TIR-domain-containing adaptor protein inducing interferon- β -mediated transcription-factor (Trif) are critical to TLR4 signaling and mediation of activation of NF- κ B. However, previous studies




Fig. 5. Effects of lumbrokinase on MPO expression in rats subjected to myocardial I-R injury. Rats were treated with vehicle or 10 $\mu\text{g}/\text{kg}$ lumbrokinase (LK-10). (A) Representative images of the Western blot results MPO expression in heart tissue samples from rats after myocardial I-R injury. (B) Quantitative densitometric analysis of MPO protein with β -actin as an internal standard. Data were normalized with sham and presented as percentage rates. Each value represents the mean of three individual experiments. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with sham. # $p < 0.05$ compared with vehicle. (C–F) Representative areas of MPO immunohistochemical (IHC) staining slides of rats subjected to myocardial I-R injury. Panels C and D are original magnification and panels E and F are higher magnification of small box in panels C and D. The red arrow indicates the infiltration of neutrophils. (G) Quantitative analysis of MPO positive cells of rat heart slides which subjected to myocardial I-R injury. Results are expressed as mean \pm SEM ($n = 3$). * $p < 0.05$ compared with vehicle.

times daily for 30 consecutive days improves regional myocardial perfusion in patients with stable angina [16]. In this study, we conducted experiments to elucidate the effectiveness of various doses of lumbrokinase. We found that 10 $\mu\text{g}/\text{kg}$ (100 units) lumbrokinase protects the heart against I-R injury in rats. In addition, we demonstrated that 10 $\mu\text{g}/\text{kg}$ lumbrokinase significantly decreases myocardial I-R-induced number of VPC and durations of VT and VF in rats. The effective dose for ameliorating myocardial I-R injury in rats is lower than that used in the clinical treatment of cardiovascular diseases.

Varda-Bloom et al. reported that immune-mediated injury from ischemia indicates that lymphocyte proliferation increases in the second week following myocardial infarction [42]. Activation of lymphocytes has been reported in later stages. In this study, we occluded the coronary artery for 1 h and then subjected it to reperfusion for 3 h. We found a significant difference in TLR4 expression in the peri-infarct area and no difference in the blood at 3 h following reperfusion. Fallach et al. also reported that they did not detect a significant infiltration of leukocytes into the heart in the acute phase (4 h), which corresponded to maximal depression of heart contraction and absence of leukocyte infiltration in TLR4-ko mice at longer time points after the LPS challenge and MI [24]. However, 24 h following MI massive leukocyte infiltration was detected. Ao et al. using isolated hearts, demonstrated that myocardial tissue TLR4, rather than neutrophil TLR4, is the determinant of myocardial neutrophil infiltration after global I-R [28]. Therefore, we suggest that TLR4 is expressed in cardiomyocytes, but not in lymphocytes, following acute myocardial I-R injury.

We also evaluated the blood-clot lysis activity of lumbrokinase. Supplementary Fig. 2 shows that treatment with 96,000 units of

lumbrokinase significantly lyses blood clots in vitro, which is consistent with the findings of a previous study [43]. However, the blood-clot lysis effect of lumbrokinase at a dose of 100 units was similar to that of vehicle. At the effective dose of lumbrokinase in rats subjected to myocardial I-R injury in this study blood-clot lysis was rarely observed. Therefore, the cardioprotective effect of lower dose lumbrokinase may be through anti-inflammatory mechanisms that protect the heart against I-R injury.

In conclusion, this study is the first to report that lumbrokinase is a potent cardioprotective drug in rats with I-R injury. Lumbrokinase decreases the number of VPC and the durations of VT and VF following I-R. We provide evidence that lumbrokinase significantly improves ventricular arrhythmias and reduces infarct volume from myocardial I-R injury. This may correlate with the inhibitory effect of the myocardial I-R-induced expressions of COX-2, iNOS, and activation of MMP-9, mediated by TLR4 signaling through JNK and NF- κ B pathways. As lumbrokinase has recently become available in tablet form and has been recommended as a dietary supplement, it is important to determine the novel signaling mechanism involved in the cardioprotective effect of lumbrokinase pretreatment. This mechanism controls innate immune response with anti-inflammatory action during myocardial I-R injury. We further conducted experiments to test the post-treatment effects of lumbrokinase in rats subjected to myocardial I-R injury. We found that 10 $\mu\text{g}/\text{kg}$ lumbrokinase significantly reduces ventricular arrhythmias and myocardial infarction in rats after myocardial I-R injury. Further studies to determine the underlying mechanisms of post-treatment lumbrokinase in the amelioration of myocardial I-R injury in rats will be undertaken. Lumbrokinase exhibits significant potential as a cardioprotective agent. Lumbrokinase may also be beneficial for

future use as a novel agent for cardiovascular disorders, such as myocardial infarction. Elucidating the mechanism by which lumbrokinase modulates the expressions of inflammatory mediators would provide a molecular basis for developing therapeutic compounds to ameliorate myocardial I-R injury.

Conflict of interest

None of the authors have any competing financial interests to declare.

Acknowledgements

This work was supported financially by research grants from Chung Shan Medical University Hospital (CSMU-INT-102-06), and the Ministry of Science and Technology (MOST 103-2320-B-040-001-MY3) of Taiwan to Dr. Shiang-Suo Huang.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.yjmcc.2016.08.004>.

References

- [1] L.M. Buja, Myocardial ischemia and reperfusion injury, *Cardiovasc. Pathol.* 14 (2005) 170–175.
- [2] M.S. Mozaffari, J.Y. Liu, W. Abebe, B. Baban, Mechanisms of load dependency of myocardial ischemia reperfusion injury, *Am J Cardiovasc Dis.* 3 (2013) 180–196.
- [3] Z.V. Schofield, T.M. Woodruff, R. Halai, M.C. Wu, M.A. Cooper, Neutrophils—a key component of ischemia-reperfusion injury, *Shock (Augusta, Ga)*, 40, 2013, pp. 463–470.
- [4] D.M. Yellon, D.J. Hausenloy, Myocardial reperfusion injury, *N. Engl. J. Med.* 357 (2007) 1121–1135.
- [5] A. Folino, G. Losano, R. Rastaldo, Balance of nitric oxide and reactive oxygen species in myocardial reperfusion injury and protection, *J. Cardiovasc. Pharmacol.* 62 (2013) 567–575.
- [6] T.V. Arumugam, I.A. Shiels, T.M. Woodruff, D.N. Granger, S.M. Taylor, The role of the complement system in ischemia-reperfusion injury, *Shock (Augusta, Ga)*, 21, 2004, pp. 401–409.
- [7] W.B. Gorsuch, E. Chrysanthou, W.J. Schwaible, G.L. Stahl, The complement system in ischemia-reperfusion injuries, *Immunobiology* 217 (2012) 1026–1033.
- [8] T.V. Arumugam, E. Okun, S.C. Tang, J. Thundyil, S.M. Taylor, T.M. Woodruff, Toll-like receptors in ischemia-reperfusion injury, *Shock (Augusta, Ga)*, 32, 2009, pp. 4–16.
- [9] H.K. Eltzschig, T. Eckle, Ischemia and reperfusion—from mechanism to translation, *Nat. Med.* 17 (2011) 1391–1401.
- [10] H. Mihara, H. Sumi, T. Yoneta, H. Mizumoto, R. Ikeda, M. Seiki, et al., A novel fibrinolytic enzyme extracted from the earthworm, *Lumbricus rubellus*, *The Japanese journal of physiology*, 41 (1991) 461–472.
- [11] I.H. Cho, E.S. Choi, H.G. Lim, H.H. Lee, Purification and characterization of six fibrinolytic serine-proteases from earthworm *Lumbricus rubellus*, *J. Biochem. Mol. Biol.* 37 (2004) 199–205.
- [12] E.L. Cooper, M. Balamurugan, Unearthing a source of medicinal molecules, *Drug Discov. Today* 15 (2010) 966–972.
- [13] R. Hu, S. Zhang, H. Liang, N. Li, C. Tu, Codon optimization, expression, and characterization of recombinant lumbrokinase in goat milk, *Protein Expr. Purif.* 37 (2004) 83–88.
- [14] L. Jin, H. Jin, G. Zhang, G. Xu, Changes in coagulation and tissue plasminogen activator after the treatment of cerebral infarction with lumbrokinase, *Clin. Hemorheol. Microcirc.* 23 (2000) 213–218.
- [15] Y.J. Cao, X. Zhang, W.H. Wang, W.Q. Zhai, J.F. Qian, J.S. Wang, et al., Oral fibrinogen-depleting agent lumbrokinase for secondary ischemic stroke prevention: results from a multicenter, randomized, parallel-group and controlled clinical trial, *Chin. Med. J.* 126 (2013) 4060–4065.
- [16] M. Kasim, A.A. Kiat, M.S. Rohman, Y. Hanifah, H. Kiat, Improved myocardial perfusion in stable angina pectoris by oral lumbrokinase: a pilot study, *J. Altern. Complement. Med.* 15 (2009) 539–544 (New York, NY).
- [17] T. Shishido, N. Nozaki, S. Yamaguchi, Y. Shibata, J. Nitobe, T. Miyamoto, et al., Toll-like receptor-2 modulates ventricular remodeling after myocardial infarction, *Circulation* 108 (2003) 2905–2910.
- [18] J. Oyama, C. Blais Jr., X. Liu, M. Pu, L. Kobzik, R.A. Kelly, et al., Reduced myocardial ischemia-reperfusion injury in toll-like receptor 4-deficient mice, *Circulation* 109 (2004) 784–789.
- [19] T. Ha, L. Liu, J. Kelley, R. Kao, D. Williams, C. Li, Toll-like receptors: new players in myocardial ischemia/reperfusion injury, *Antioxid. Redox Signal.* 15 (2011) 1875–1893.
- [20] Liao PH, Hung LM, Chen YH, Kuan YH, Zhang FB, Lin RH, et al. Cardioprotective effects of luteolin during ischemia-reperfusion injury in rats. *Circ. J. (official journal of the Japanese Circulation Society)*. 2011;75:443–50.
- [21] M.J. Curtis, J.C. Hancox, A. Farkas, C.L. Wainwright, C.L. Stables, D.A. Saint, et al., The Lambeth conventions (II): guidelines for the study of animal and human ventricular and supraventricular arrhythmias, *Pharmacol. Ther.* 139 (2013) 213–248.
- [22] M.J. Walker, M.J. Curtis, D.J. Hearse, R.W. Campbell, M.J. Janse, D.M. Yellon, et al., The Lambeth conventions: guidelines for the study of arrhythmias in ischaemia infarction, and reperfusion, *Cardiovasc. Res.* 22 (1988) 447–455.
- [23] E. Matsuura, F. Umehara, T. Hashiguchi, N. Fujimoto, Y. Okada, M. Osame, Marked increase of matrix metalloproteinase 9 in cerebrospinal fluid of patients with fungal or tuberculous meningoencephalitis, *J. Neurol. Sci.* 173 (2000) 45–52.
- [24] R. Fallach, A. Shainberg, O. Avlas, M. Fainblut, Y. Chepurko, E. Porat, E. Hochhauser, Cardiomyocyte toll-like receptor 4 is involved in heart dysfunction following septic shock or myocardial ischemia, *J. Mol. Cell. Cardiol.* 48 (2010) 1236–1244.
- [25] O. Avlas, R. Fallach, A. Shainberg, E. Porat, E. Hochhauser, Toll-like receptor 4 stimulation initiates an inflammatory response that decreases cardiomyocyte contractility, *Antioxid. Redox Signal.* 15 (2011) 1895–1909.
- [26] P. Zhao, J. Wang, L. He, H. Ma, X. Zhang, X. Zhu, et al., Deficiency in TLR4 signal transduction ameliorates cardiac injury and cardiomyocyte contractile dysfunction during ischemia, *J. Cell. Mol. Med.* 13 (2009) 1513–1525.
- [27] A.K. Chow, J. Cena, R. Schulz, Acute actions and novel targets of matrix metalloproteinases in the heart and vasculature, *Br. J. Pharmacol.* 152 (2007) 189–205.
- [28] L. Ao, N. Zou, J.C. Cleveland Jr., D.A. Fullerton, X. Meng, Myocardial TLR4 is a determinant of neutrophil infiltration after global myocardial ischemia: mediating KC and MCP-1 expression induced by extracellular HSC70, *Am. J. Physiol. Heart Circ. Physiol.* 297 (2009) H21–H28.
- [29] J.H. Boyd, S. Mathur, Y. Wang, R.M. Bateman, K.R. Walley, Toll-like receptor stimulation in cardiomyocytes decreases contractility and initiates an NF-kappaB dependent inflammatory response, *Cardiovasc. Res.* 72 (2006) 384–393.
- [30] E. Wang, Y. Feng, M. Zhang, L. Zou, Y. Li, E.S. Buys, et al., Toll-like receptor 4 signaling confers cardiac protection against ischemic injury via inducible nitric oxide synthase- and soluble guanylate cyclase-dependent mechanisms, *Anesthesiology* 114 (2011) 603–613.
- [31] Y. Li, R. Si, Y. Feng, H.H. Chen, L. Zou, E. Wang, et al., Myocardial ischemia activates an injurious innate immune signaling via cardiac heat shock protein 60 and toll-like receptor 4, *J. Biol. Chem.* 286 (2011) 31308–31319.
- [32] F. Hua, T. Ha, J. Ma, X. Gao, J. Kelley, D.L. Williams, et al., Blocking the MyD88-dependent pathway protects the myocardium from ischemia/reperfusion injury in rat hearts, *Biochem. Biophys. Res. Commun.* 338 (2005) 1118–1125.
- [33] Q. Jiang, M. Yi, Q. Guo, C. Wang, H. Wang, S. Meng, et al., Protective effects of polydatin on lipopolysaccharide-induced acute lung injury through TLR4-MyD88-NF-kappaB pathway, *Int. Immunopharmacol.* (2015).
- [34] Y. Fang, J. Hu, Toll-like receptor and its roles in myocardial ischemic/reperfusion injury, *Med. Sci. Monit.* 17 (2011) Ra100–9 (international medical journal of experimental and clinical research).
- [35] M. Sato, D. Bagchi, A. Tosaki, D.K. Das, Grape seed proanthocyanidin reduces cardiomyocyte apoptosis by inhibiting ischemia/reperfusion-induced activation of JNK-1 and C-JUN, *Free Radic. Biol. Med.* 31 (2001) 729–737.
- [36] G. Vassalli, G. Milano, T. Moccetti, Role of mitogen-activated protein kinases in myocardial ischemia-reperfusion injury during heart transplantation, *J. Transplant.* 2012 (2012) 928954.
- [37] K.K. Wu, Inducible cyclooxygenase and nitric oxide synthase, *Adv. Pharmacol.* 33 (1995) 179–207 (San Diego, Calif).
- [38] M. Lindsey, K. Wedin, M.D. Brown, C. Keller, A.J. Evans, J. Smolen, et al., Matrix-dependent mechanism of neutrophil-mediated release and activation of matrix metalloproteinase 9 in myocardial ischemia/reperfusion, *Circulation* 103 (2001) 2181–2187.
- [39] A.M. Romanic, S.M. Harrison, W. Bao, C.L. Burns-Kurtis, S. Pickering, J. Gu, et al., Myocardial protection from ischemia/reperfusion injury by targeted deletion of matrix metalloproteinase-9, *Cardiovasc. Res.* 54 (2002) 549–558.
- [40] L. Lu, Z. Gunja-Smith, J.F. Woessner, P.C. Ursell, T. Nissen, R.E. Galaray, et al., Matrix metalloproteinases and collagen ultrastructure in moderate myocardial ischemia and reperfusion in vivo, *Am. J. Physiol. Heart Circ. Physiol.* 279 (2000) H601–H609.
- [41] M.K. Verma, K.K. Pulicherla, Lumbrokinase – a potent and stable fibrin-specific plasminogen activator, *Int. J. Biosci. Biotechnol.* 3 (2011) 57–70.
- [42] N. Varda-Bloom, J. Leor, D.G. Ohad, Y. Hasin, M. Amar, R. Fixler, et al., Cytotoxic T lymphocytes are activated following myocardial infarction and can recognize and kill healthy myocytes in vitro, *J. Mol. Cell. Cardiol.* 32 (2000) 2141–2149.
- [43] G. Li, K.Y. Wang, D. Li, N. Wang, D. Liu, Cloning, expression and characterization of a gene from earthworm *Eisenia fetida* encoding a blood-clot dissolving protein, *PLoS One* 7 (2012), e31110.

103年度專題研究計畫成果彙整表

計畫主持人：黃相碩			計畫編號：103-2320-B-040-001-MY3				
計畫名稱：抗發炎機制在於Lumbrokinase心臟保護作用中扮演的角色							
成果項目			量化	單位	質化 (說明：各成果項目請附佐證資料或細項說明，如期刊名稱、年份、卷期、起訖頁數、證號...等)		
國內	學術性論文	期刊論文		0	篇		
		研討會論文		0			
		專書		0	本		
		專書論文		0	章		
		技術報告		0	篇		
		其他		0	篇		
	智慧財產權及成果	專利權	發明專利	申請中	0	件	
				已獲得	0		
			新型/設計專利		0		
		商標權		0			
		營業秘密		0			
		積體電路電路布局權		0			
		著作權		0			
		品種權		0			
		其他		0			
	技術移轉	件數		0	件		
		收入		0	千元		
	國外	學術性論文	期刊論文		1	篇	Journal of Molecular and Cellular Cardiology 99 (2016) 113 - 122
			研討會論文		0		
			專書		0	本	
專書論文			0	章			
技術報告			0	篇			
其他			0	篇			
智慧財產權及成果		專利權	發明專利	申請中	0	件	
				已獲得	0		
			新型/設計專利		0		
		商標權		0			
		營業秘密		0			
		積體電路電路布局權		0			
		著作權		0			
		品種權		0			

		其他	0		
	技術移轉	件數	0	件	
		收入	0	千元	
參與計畫人力	本國籍	大專生	4	人次	蔡仁維；周嘉宇；張家崧；姚懿庭
		碩士生	1		洪岳慈
		博士生	2		王羿忻；蘇偉慧
		博士後研究員	0		
		專任助理	0		
	非本國籍	大專生	0		
		碩士生	0		
		博士生	0		
		博士後研究員	0		
		專任助理	0		
其他成果 (無法以量化表達之成果如辦理學術活動、獲得獎項、重要國際合作、研究成果國際影響力及其他協助產業技術發展之具體效益事項等，請以文字敘述填列。)					

科技部補助專題研究計畫成果自評表

請就研究內容與原計畫相符程度、達成預期目標情況、研究成果之學術或應用價值（簡要敘述成果所代表之意義、價值、影響或進一步發展之可能性）、是否適合在學術期刊發表或申請專利、主要發現（簡要敘述成果是否具有政策應用參考價值及具影響公共利益之重大發現）或其他有關價值等，作一綜合評估。

1. 請就研究內容與原計畫相符程度、達成預期目標情況作一綜合評估

達成目標

未達成目標（請說明，以100字為限）

實驗失敗

因故實驗中斷

其他原因

說明：

2. 研究成果在學術期刊發表或申請專利等情形（請於其他欄註明專利及技轉之證號、合約、申請及洽談等詳細資訊）

論文： 已發表 未發表之文稿 撰寫中 無

專利： 已獲得 申請中 無

技轉： 已技轉 洽談中 無

其他：（以200字為限）

3. 請依學術成就、技術創新、社會影響等方面，評估研究成果之學術或應用價值（簡要敘述成果所代表之意義、價值、影響或進一步發展之可能性，以500字為限）

心肌缺血再灌注傷害是冠狀動脈疾病發生以及死亡的主要原因。本研究在大白鼠左冠狀動脈下降支結紮一小時而後再灌注三小時之心肌缺血再灌注傷害動物模式，發現lumbrokinase會減少心肌缺血再灌注傷害造成之動物死亡率以及心肌缺血再灌注傷害所誘發之心律不整，投予lumbrokinase會減少心肌缺血再灌注傷害所造成的心肌梗塞的體積，在心肌缺血再灌注傷害期間

，lumbrokinase可以改善心臟收縮與舒張功能，並降低動脈血漿中心肌細胞傷害因子LDH的活性，Lumbrokinase經由TLR4訊息傳遞路徑抑制心肌缺血再灌注損傷增加的COX-2，iNOS以及MMP-9的蛋白質表現。此外，我們的研究結果證明，Lumbrokinase減少JNK，IkB和NF- κ B的磷酸化。這些結果可以證明

lumbrokinase是一個很有潛力的心臟保護劑，可以保護心臟對抗心肌缺血再灌注傷害，具有抗心律不整與減少心肌梗塞的心臟保護作用。此研究成果已發表於Journal of Molecular and Cellular Cardiology 2016; 99: 113 - 122。

(IF=5.68)

4. 主要發現

本研究具有政策應用參考價值：否 是，建議提供機關
(勾選「是」者，請列舉建議可提供施政參考之業務主管機關)

本研究具影響公共利益之重大發現：否 是

說明：(以150字為限)