

## 參考文獻

- Andrew, S. E., Goldberg Y. P., Kremer, B., Telenius, H., Theimann, J., Adam, S., Starr, E., Squitieri, F., Lin, B., Kalchman, M. A., Graham, R. K., and Hayden, M. R. (1993) The relationship between trinucleotide (CAG) repeat length and clinical features of Huntington's disease. *Nature Genet.* **4**, 398-403.
- Amack JD, Paguio AP, Mahadevan MS. (1999) Cis and trans effects of the myotonic dystrophy (DM) mutation in a cell culture model. *Hum Mol Genet.* **8**(11), 1975-84.
- Amack JD, Mahadevan MS. (2001) The myotonic dystrophy expanded CUG repeat tract is necessary but not sufficient to disrupt C2C12 myoblast differentiation. *Hum Mol Genet.* **10**(18), 1879-87.
- Amack JD, Reagan SR, Mahadevan MS. (2002) Mutant DMPK 3'-UTR transcripts disrupt C2C12 myogenic differentiation by compromising MyoD. *J Cell Biol.* **159**(3), 419-29.
- Ashizawa, T. and Epstein, H. F. (1991) Ethnic distribution of the myotonic dystrophy gene. *Lancet* **338**, 642-643.
- Aslanidis, C., Jansen, G., Amemiya, C., Shutler, G., Mahadevan, M., Tsilfidis, C. et al. (1992) Cloning of the essential myotonic dystrophy region: mapping of the putative defect. *Nature* **355**, 548-551.
- Berul, C. I., Maguire, C. T., Aronovitz, M. J., Greenwood, J., Miller, C., Gehrmann, J., Housman, D., Mendelsohn, M. E., and Reddy, S. (1999) DMPK dosage alternations result in atrioventricular conduction abnormalities in a mouse myotonic dystrophy model. *J. Clin. Invest.* **103**(4), R1-7.

- Bouchard, G., Roy, R., Declos, M., Mathieu, J., and Kouladjian, K. (1989) Origin and diffusion of the myotonic dystrophy gene in the Saguenay region (Quebec). *Can. J. Neurol Sci.* **116**, 119-122.
- Boucher, C. A., King, S. K., Carey, N., Krahe, R., Winchester, C. L., Rahman, S., Creavin, T., Meghji, P., Bailey M. E. S., Chartier, F. L., Brown, S. D., Siciliano, M. J., and Johnson, K. L. (1995) A novel homeodomain-encoding gene is associated with a large CpG island interrupted by the myotonic dystrophy unstable (CTG)<sub>n</sub> repeat. *Hum. Mol. Genet.* **4**, 1919-1925.
- Brook, J. D., McCurrach, M. E., Harley, H. G., Buckler, A. J., Church, D., Aburatani, H., Hunter, K., Stanton, V. P., Thirion, J.-P., Hudson, T., Sohn, R., Zemelmann, B., Snell, R. G., Rundle, S. A., Crow, S., Davies, J., Shelbourne, P., Buxton, J., Johns, C., Juvonen, V., Johnson, K., Harper, P. S., Shaw, D. J., and Housman D. E. (1992) Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* **68**, 799-808.
- Buxton, J., Schelbourne, P., Davies, J., Jones, C., Van Tongeren, T., Aslanidis, C., De Jong, P., Jansen, G., Anvert, M. et al.. (1992) Detection of an unstable fragment of DNA specific to individuals with myotonic dystrophy. *Nature* **355**, 547-548.
- Charlet-B N, Savkur RS, Singh G, Philips AV, Grice EA, Cooper TA. (2002) Loss of the muscle-specific chloride channel in type 1 myotonic dystrophy due to misregulated alternative splicing. *Mol Cell*.**10**(1), 45-53.
- Davies, J., Yamagata, H., Shelbourne, P., Buxton, J., Ogihara, T.,

- Nokelainen, P., Nakagawa, M., et al. (1992) Comparison of the myotonic dystrophy associated CTG repeat in European and Japanese populations. *J. Med. Genet.* **29**, 766-769.
- Davis, B. M., McCurrach, M. E., Taneja, K. L., Singer, R. H., and Housman, D. E. (1997) Expansion of a CUG trinucleotide repeat in the 3' untranslated region of myotonic dystrophy protein kinase transcripts results in nuclear retention of transcripts. *Proc. Natl. Acad. Sci., USA* **94** (14), 7388-7393.
- Duyao, M., Ambrose, C., Myers, R., Novelletto, A., Persichetti, F., Fontali, M., Folstein, S. et al. (1993) Trinucleotide repeat length instability and age of onset in Huntington's disease. *Nature Genet.* **4**, 387-392.
- Emmanuelle Fabre, Bernard Dujon and Guy-Franck Richard (2002) Transcription and nuclear transport of CAG/CTG trinucleotide repeats in yeast. *Nucleic Acids Res.* **30**(16):3540-7.
- Fu, Y.-H., Friedman, D. L., Richards, S., Pearlman, J. A., Gibbs, R. A., Pizzuti, A., Ashizawa, T., Perryman, M. B., Scariato, G., Fenwick, R. G., Jr., and Caskey, C. T. (1993) Decreased expression of myotonin-protein kinase messenger RNA and protein in adult form of myotonic dystrophy. *Science* **260**, 235-238.
- Fu, Y.-H., Pizzuti, A., Fenwick, R. G., King, J., Jr., Pajnarayan, S., Dunne, P. W., Dubel, J., Nasser, G. A., Ashizawa, T., de Jong, P., Wieringa, B., Korneluk, R., Perryman, M. B., Epstein, H. F., and Caskey, C. T. (1992) An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science* **255**, 1256-1258.
- George, Jr. A. L., Crackower, M. A., Abdalla, J. A., Hudson, A. J., and

- Ebers, G. C. (1993) Molecular basis of Thomsen's disease (autosomal dominant myotonia congenital). *Nature Genet.* **3**, 305-310.
- Harley, H. G., Brook, J. D., Rundle, S. A., Crow, S., Reardon, W., Buckler, A. J., Harper, P. S., Housman, D. E., and Shaw, D. J. (1992) Expansion of an unstable DNA region and phenotypic variation in myotonic dystrophy. *Nature* **355**, 545-546.
- Harper P. S. (1989) Myotonic dystrophy, 2<sup>nd</sup> ed., WB Saunders, London.
- Harris, S., Moncrieff, C., and Johnson, K. (1996) Myotonic dystrophy: will the real gene please step forward! *Hum. Mol. Genet.* **5**, 1417-1423.
- Heath, S. K., Carne, S., Hoyle, C., Johnson, K. L., and Wells, D. J. (1997) Characterization of expression of mDMAHP, a homeodomain-encoding gene at the murine DM locus. *Hum. Mol. Genet.* **6**, 651-657.
- Seznec H, Agbulut O, Sergeant N, Savouret C, Ghestem A, Tabti N, Willer JC, Ourth L, Duros C, Brisson E, Fouquet C, Butler-Browne G, Delacourte A, Junien C, Gourdon G. (2001) Mice transgenic for the human myotonic dystrophy region with expanded CTG repeats display muscular and brain abnormalities. *Hum. Mol. Genet.* **10**, 2717-2726.
- Hiroyuki, K., Takahisa, T., Kwasi, G. M., Akiko, W., Masato, T., and Toshiaki, I. (2000) Profiling of *Caenorhabditis elegans* proteins using two-dimensional gel electrophoresis and matrix assisted laser desorption/ionization-time of flight-mass spectrometry. *Electrophoresis* **21**, 1755-1765.
- Hoffman-Radvanyi, H., Lavedan, H., Rabes, J. P., Savoy, D., Duros, C.,

- Johnson, K., and Junien, K. (1993) Myotonic dystrophy: absence of CTG enlarged transcript in congenital forms, and low expression of the normal allele. *Hum. Mol. Genet.* **2**, 1263-1266.
- Hsiao, K.-M., Chou, S.-Y., Shih, S.-J., and Ferrell, J. E., Jr. (1994) Evidence that inactive p42 mitogen-activated protein kinase and inactive Rsk exist as a heterodimer in vivo. *Proc. Natl. Acad. Sci. USA* **91**, 5480-5484.
- Hsiao, K.-M., McMahon, S. L., and Farnham, P. J. (1994) Multiple DNA elements are required for the growth regulation of the mouse E2F1 promoter. *Genes & Dev.* **8**, 1526-1537.
- Hsiao, K.-M., Lin, H.-M., Pan, H., Li, T.-C., Chen, S.-S., Jou, S.-B., Chiu, Y.-L., Wu, M.-F., Lin, C.-C., and Li, S.-Y. (1999) Application of FTA sample collection and DNA purification system on the determination of CTG trinucleotide repeat size by PCR-based Southern blotting. *J. Clin. Lab. Anal.* **13**, 188-193.
- Hsieh, M., Lin, S. J., Chen, J. F., Lin, H. M., Hsiao, K. M., Li, S. Y., Li, C., and Tsai, C. J. (2000) Identification of the spinocerebellar ataxia type 7 mutation in Taiwan: application of PCR-based Southern blot. *J. Neurol.* **247**, 623-629.
- Jansen, G., Groenen, P. J., T. A., Bachner, D., Jap, P. H. K., Coerwinkel, M., Oerlemans, F., van den Broek, W., Gohlsch, B., Pette, D., Plomp, J. J., Molenaar, P. C., Nederhoff, M. G. J., van Echteld, C. J. A., Dekker, M., Berns, A., Hameister, H., and Wieronga, B. (1996) Abnormal myotonic dystrophy protein kinase levels produce only mild myopathy in mice. *Nature Genet.* **13**, 316-324.
- Jodice, C., Malaspina, P., Persichetti, F., Novelletto, A., Spadaro, M.,

- Giunti, P., Morocutti, C. et al. (1994) Effect of trinucleotide repeat length and parental sex on phenotypic variation in spinocerebellar ataxia 1. *Am. J. Hum. Genet.* **54**, 959-965.
- Jou, S.-B., Lin, H.-M., Pan, H., Chiu, Y.-L., Li, S.-Y., and Hsiao, K.-M. (2001). Delineation of CTG repeats and clinical features in Taiwanese Myotonic Dystrophy family. *Proc. Natl. Sci. Council, ROC.* **25**, 48-52.
- Kawaguchi, Y., Okamoto, T., Taniwaki, M., Aizawa, M., Inoue, M., Katayama, S., kawakami, H. et al. (1994) CAG expansions in a novel gene from Machado-Joseph Disease at chromosome 14q32.1. *Nature Genet.* **8**, 221-227.
- Klesert, T. R., Otten, A. D., Bird, T. D., and Tapscott, S. J. (1997) Trinucleotide repeat expansion at the myotonic dystrophy locus reduces expression of DMAHP. *Nature Genet.* **16**, 402-406.
- Klesert, T. R., Cho, D. H., Clark, J. I., Maylie, J., Adelman, J., Snider, L., Yuen, E. C., Soriano, P., and Tapscott, S. J. (2000) Mice deficient in Six5 develop cataracts: implications for myotonic dystrophy. *Nature Genet.* **25**, 105-109.
- Koide, R., Ikeuchi, T., Onodera, O., Tanaka, H., Igarashi, S., Endo, K., Takahashi, H., Kondo, R., Ishiawa, A., Hayashi, T., Saito, M., Tomoda, A., Miike, T., Naito, H., Ikuta, F., and Tsuji, S. (1994) Unstable expansion of CAG repeat in hereditary dentatorubral-pallidoluysian atrophy (DRPLA). *Nature Genet.* **6**, 9-13.
- Korade-Mirnic, Z., Bantzke, P., and Hoffman, E. (1998) Myotonic dystrophy: molecular windows on a complex etiology. *Nucleic Acids Res.* **26**, 1363-1368.

- La Spada. A. R., Wilson, E. M., Lubahn, D. B., Harding, A. E., and Fischbeck, K. H. (1991) Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* **352**, 77-79.
- Lin, M. J. and Lin-Shiau, S. Y. (2000) Suramin protects the murine motor nerves from the toxic effects of presynaptic  $\text{Ca}^{2+}$  channel inhibitors. *Neurosci. Letters* **287**, 97-100.
- Lin-Shiau, S. Y. and Lin, M. J. (1999) Suramin inhibits the toxic effects of presynaptic neurotoxins at the mouse motor nerve terminals. *Euro. J. Pharmacol.* **382**, 75-80.
- Lin, M. J. and Lin-Shiau, S. Y. (1999) Enhanced spontaneous transmitter release at murine motor nerve terminals with cyclosporine. *Neuropharmacol.* **38**, 195-198.
- Lin, M. J. and Lin-Shiau, S. Y. (1998) Phorbol ester-induced contractions of mouse detrusor muscle are inhibited by nifedipine. *Naunyn-Schmiedeberg's Arch Pharmacol.* **357**, 553-557.
- Lin-Shiau, S. Y. and Lin, M. J. (1998) Studies on curare-like action of the tripeptide carbobenzoxy-Gly-Gly-Arg- -naphthylamide in mouse diaphragm. *Euro. J. Pharmacol.* **343**, 51-56 .
- Lin, M. J. and Lin-Shiau, S. Y. (1997) Multiple types of  $\text{Ca}^{2+}$  channels at the mouse motor nerve terminals. *Euro. J. Neurosci.* **9**: 817-823 (1997).
- Lin, M. J. and Lin-Shiau, S. Y. (1996) Ruthenium Red, a novel enhancer of  $\text{K}^{+}$  currents at mouse motor nerve terminals. *Neuropharmacol.* **35**, 615-623.
- Lin, M. J. and Lin-Shiau, S. Y. (1995) Functional assessment of  $\text{Ca}^{2+}$ -current in the mouse motor nerve terminals. *Neurosci. Letter* **195**,

21-24.

Lin-Shiau, S. Y., Liu, S. H., and Lin, M.J. (1993) Use of ion channel blockers in the exploration of possible mechanisms involved in the myopathy of diabetic mice. *Naunyn-Schmiedeberg Arch. Pharmacol.* **348**: 311-318.

Maciel, P., Gaspar, C., DeStefano, A. L., Silveira, L., Coutinho, P., Radvany, J., Dawson, D. M., Sundarsky, L., Guimaraes, J., Loureiro, J. E. L., Nezarati, M. M., Corwin, L. L., Lopes-Cendes, I., Rooke, K., Rosenberg, R., MacLeod, P., Farrer, L. A., Sequeiros, J., and Rouleau, G. A. (1995) Correlation between CAG repeat length and clinical features in Machado-Joseph disease. *Am. J. Hum. Genet.* **57**, 54-61.

Mahadevan, M., Tsilfidis, C., sabourin, L., Shutler, G., Amemiya, C., Jansen, G., Neville, C., Narang, M., Barcero, J., O'Hoy, K., Leblond, S., Earle-Macdonald, J., de Jong, P. J., Wieringa, B., and Korneluk, R. G. (1992) Myotonic dystrophy mutation: an unstable CTG repeat in the 3' untranslated region of the gene. *Science* **255**, 1253-1255.

Mankodi, A., Logigian, E., Callahan, L., McClain, C., White, R., Henderson, D., Krym, M., and Thornton, C., A. (2000) Myotonic dystrophy in transgenic mice expressing an expanded CUG repeat. *Science* **289**, 1769-1772.

Miller JW, Urbinati CR, Teng-Umnuay P, Stenberg MG, Byrne BJ, Thornton CA, Swanson MS. (2000) Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J*, **19**(17) 4439-48.

Fardaei M, Rogers MT, Thorpe HM, Larkin K, Hamshire MG, Harper PS, Brook JD. (2002) Three proteins, MBNL, MBLL and MBXL,



- co-localize in vivo with nuclear foci of expanded-repeat transcripts in DM1 and DM2 cells. *Hum Mol Genet.* **11**(7), 805-14.
- Fardaei M, Larkin K, Brook JD, Hamshere MG. (2001) In vivo co-localisation of MBNL protein with DMPK expanded-repeat transcripts. *Nucleic Acids Res.* 2001 **29**(13), 2766-71.
- Mankodi A, Takahashi MP, Jiang H, Beck CL, Bowers WJ, Moxley RT, Cannon SC, Thornton CA. (2002) Expanded CUG repeats trigger aberrant splicing of ClC-1 chloride channel pre-mRNA and hyperexcitability of skeletal muscle in myotonic dystrophy. *Mol Cell.***10**(1):35-44
- Milne, C. A., and Hodgkin, J. (1999) ETR-1, a homologue of a protein linked to myotonic dystrophy, is essential for muscle development in *Caenorhabditis elegans*. *Current Biology* **9**, 1243-1246.
- Morrone, A., Pegoraro, E., Angelini, C., Zammarchi, E., Marconi, G., and Hoffman, E. (1997) RNA metabolism in myotonic dystrophy: patient muscle shows decreased insulin receptor RNA and protein consistent with abnormal insulin resistance. *J. Clin. Invest.* **99**, 1691-1698.
- Nelson, D. L. (1996) Allelic expansion underlies many genetic diseases. *Growth: Genetics & Hormones* **12**, 1-4.
- Noguchi S, Wakabayashi-Takai E, Sasaoka T, Ozawa E. (2001) Analysis of the spatial, temporal and tissue-specific transcription of gamma-sarcoglycan gene using a transgenic mouse. *FEBS Lett.* **495**(1-2), 77-81.
- Otten, A.D. and Tapscott, S. J. (1995) Triplet repeat expansion in myotonic dystrophy alters the adjacent chromatin structure. *Proc. Natl. Acad. Sci. USA* **92**, 5465-5469.

- Pan, H. and Griep, A. E. (1994) Altered cell cycle regulation in the lens of HPV-16 E6 or E7 transgenic mice: implications of tumor suppressor gene function in development. *Genes & Dev.* **8**, 1285-1299.
- Pan, H. and Griep, A. E. (1995) Temporally distinct patterns of p53-dependent and p53-independent apoptosis during mouse lens development. *Genes & Dev.* **9**, 2157-2169.
- Pan, H., Yin, C., and Van Dyke T. A. (1997) Apoptosis and cancer mechanisms. *Cancer Surveys* **29**, 305-327.
- Pan, H., Yin, C., Dyson, N.J., Harlow, E., Yamasaki, L., and Van Dyke, T. (1998) Key roles for E2F1 in signaling p53-dependent apoptosis and in cell division within developing tumor. *Mol. Cell* **2**, 283-292.
- Pan, H., Lin, H.-M., Ku, W.-Y., Li, T.-C., Li, S.-Y., Lin, C. C., and Hsiao, K.-M. (submitted-1) Haplotype analysis of the myotonic dystrophy type 1 (DM1) locus in Taiwan: implications for low prevalence and founder mutations of Taiwanese myotonic dystrophy.
- Pan, H., Li, Y.-Y., Li, T.-C., Tsai, W.-T., Li, S.-Y., and Hsiao, K.-M. (submitted-2) Trinucleotide repeat lengths in the myotonic dystrophy and Machado-Joseph disease genes are enhanced in idiopathic azoospermia patients.
- Phelps, D.E., Hsiao, K.-M., Li, Y., Hu, N., Franklin, D.S., Westphal, E., Lee, E.Y.-H.P., and Xiong, Y. (1998) Coupled transcriptional and translational control of cyclin-dependent kinase inhibitor p18<sup>INK4c</sup> expression during myogenesis. *Mol. Cell Biol.* **18**, 2334-2343.
- Philips, A.V., Timchenko, L.T., and Cooper, T.A. (1998) Disruption of splicing regulated by a CUG-binding protein in myotonic dystrophy. *Science* **280**, 737-741.

- Ranum, L. P. W., Chung, M., Banfi, S., Bryer, A., Schut, L. J., Rameser, R., Duvick, L. A. et al. (1994) Molecular and clinical correlations in spinocerebellar ataxia type 1: evidence for familial effects on the age at onset. *Am. J. Hum. Genet.* **55**, 244-252.
- Ranum, L.P.W., Rasmussen, P.F., Benzow, K.A., Koob, M.D., and Day, J.W. (1998) Genetic mapping of a second myotonic dystrophy locus. *Nature Genet.* **19**, 196-198.
- Reddy, S., Smith, D. B. J., Rich, M. M., Leferovich, J. M., Reilly, P., Davis, B. M., Tran, K., rayburn, H., Bronson, R., Cros D., Balice-Gordon, R. J., and Housman, D. (1996) Mice lacking the myotonic dystrophy protein kinase develop a late onset progressive myopathy. *Nature Genet.* **13**, 325-335.
- Ricker, K., Koch, M. C., Lehmann-Horn, F. et al. (1995) Proximal myotonic myopathy. Clinical features of a multisystem disorder similar to myotonic dystrophy. *Arch. Neurol.* **52**, 25-31.
- Sabourin, L., Mahadevan, M. S., Narang, M., Lee, D. S. C., Surh, L. C., and Korneluk, R. G. (1993) Effect of the myotonic dystrophy (DM) mutation on mRNA levels of the DM gene. *Nature Genet.* **4**, 233-238.
- Savkur RS, Philips AV, Cooper TA. (2001) Aberrant regulation of insulin receptor alternative splicing is associated with insulin resistance in myotonic dystrophy. *Nat Genet.* **29**(1), 40-7.
- Sarker, P. S., Appukuttan, B., Han, J., Ito, Y., Ai, C., Tsai, W., Chai, Y., Stout, J. T., and Reddy, S. (2000) Heterozygous loss of Six5 in mice is sufficient to cause ocular cataracts. *Nature Genet.* **25**, 110-114.
- Snell, R. G., MacMillan, J. C., Cheadle, J. P., Fenton, I., Lazarou, L. P., Davies, P., MacDonald, M. E., Gusella, J. F., Harper, P. S., and Shaw,

- D. J. (1993) Relationship between trinucleotide repeat expansion and phenotypic variation in Huntington's disease. *Nature Genet.* **4**, 393-397.
- Squillace RM, Chenault DM, Wang EH. (2002) Inhibition of muscle differentiation by the novel muscleblind-related protein CHCR. *Dev Biol.* **250**(1):218-30.
- Taneja, K. L., McCurrach, M., Schalling, M., Housman, D., and Singer, R. H. (1995) Foci of trinucleotide repeat transcripts in nuclei of myotonic dystrophy cells and tissue. *J. Cell Biol.* **128**, 995-1002.
- The international Myotonic Dystrophy Consortium (IDMC) (2000) New nomenclature and DNA testing guidelines for myotonic dystrophy type 1 (DM1). *Neurology* **54**: 1218-1221.
- Thornton C. A., Wymer, J. P., Simmons, Z., McClain, C., and Moxley, R. T. (1997) Expansion of the myotonic dystrophy CTG repeat reduces expression of the flanking DMAHP gene. *Nature Genet.* **16**, 407-409.
- Timchenko, L. T., and Caskey, C. T. (1996) Trinucleotide repeat disorders in humans: discussions of mechanisms and medical issues. *FASEB J.* **10**, 1589-1597.
- Timchenko, L.T., Miller, J.W., Timchenko, N.A., De Vore, D.R., Datar, K.V., Lin, L., Roberts, R., Caskey, C.T., and Swanson, M.S. (1996a) Identification of a (CUG)<sub>n</sub> triplet repeat RNA-binding protein and its expression in myotonic dystrophy. *Nucleic Acids Res.* **24**, 4407-4414.
- Timchenko, L.T., Timchenko, N.A., Caskey, C.T., and Roberts, R. (1996b) Novel proteins with binding activity for DNA CTG and RNA CUG repeats: implications for myotonic dystrophy. *Hum. Mol. Genet.* **5**, 115-121.

- Timchenko NA, Cai ZJ, Welm AL, Reddy S, Ashizawa T, Timchenko LT. (2001) RNA CUG repeats sequester CUGBP1 and alter protein levels and activity of CUGBP1. *J Biol Chem.* **276**(11) 7820-6
- Udd, B., Krahe, R., Wallgren-Pettersson, C., Falck, B., and Kalimo, H. (1997) Proximal myotonic dystrophy- a family with autosomal dominant muscular dystrophy, cataracts, hearing loss and hypogonadism: heterogeneity of proximal myotonic syndromes? *Neuromuscul. Disord.* **7**, 217-228.
- Verkerk, A.J.M.H., Pieretti, M., Sutcliffe, J. S., Fu, Y.-H., Kuhl, D. P. A., Pizzuti, A., Reiner, O., Richards, S., Victoria, M. F., Zhang, F., Eussen, B. E., van Ommen, G.-J. B., Blonden, L. A. J., Riggins, G. J., Chastain, J. L., Kunst, C. B., Galjaard, H., caskey, C. T., Nelson, D. L., Ooster, B. A., and Warren, S. T. (1991) Identification of a gene (*FMR-1*) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* **65**, 905-914.
- VanGinkel, P.R., Hsiao, K.-M., Schjerven, H., and Farnham, P.J. (1997) E2F-mediated growth regulation requires transcription factor cooperation. *J. Bio. Chem.* **272**(29), 18367-18374.
- Verma, A. K., Hsiao, K.-M., Ahrens, H., Suganuma, M., Fujiki, H., Matsufuji, S., and Hayashi, H. (1996) Superinduction of mouse epidermal ornithine decarboxylase activity by repeated 12-o-tetradecanoylphorbol-13-acetate treatments. *Mol. Cell. Biochem.* **155**(2), 139-151.
- Vincent, J. B., Paterson, A. D., Strong, E., Petronis, A., and Kennedy, J. L. (2000) The unstable trinucleotide repeat story of major psychosis. *Am. J. Med. Genet.* **97**, 77-97.

- Wakabayashi-Takai E, Noguchi S, Ozawa E. (2001) Identification of myogenesis-dependent transcriptional enhancers in promoter region of mouse gamma-sarcoglycan gene. *Eur J Biochem.* **268**(4), 948-57.
- Wang, Y.-H., Amirhaeri, S., Kang, S., Wells, R. D., and Griffith, J. D. (1994) Preferential nucleosome assembly at DNA triplet repeats from the myotonic dystrophy gene. *Science* **265**, 669-671.
- Wang, Y.-H. and Griffith, J. (1995) Expanded CTG triplet blocks from the myotonic dystrophy gene create the strongest known natural nucleosome positioning elements. *Genomics* **25**, 570-573.
- Wissmann, A., Ingles, J., McGhee J. D., and Mains, P. E. (1997) *Caenorhabditis elegans* LET-502 is related to Rho-binding kinases and human myotonic dystrophy kinase and interacts genetically with a homolog of the regulatory subunit of smooth muscle myosin phosphatase to affect cell shape. *Genes & Dev.* **15**, 409-422.
- Wooster, R., Cleton-Jansen, A. M., Collins, N., Mangion, J., Cornelis, R. S., Cooper, C. S., Gusteson, B. A., Ponder, B. A., von Deimling, A., Wiestler, O. D., et al. (1994) Instability of short tandem repeats (microsatellites) in human cancers. *Nature Genet.* **6**, 152-156.
- Wu, Y. and Horvitz, H. R. (1998a) *C. elegans* phagocytosis and cell-migration protein CED-5 is similar to human DOCK180. *Nature* **392**, 501-504.
- Wu, Y. and Horvitz, H. R. (1998b) The *C. elegans* cell corpse engulfment gene *ced-7* encodes a protein similar to ABC transporters. *Cell* **93**, 951-960.