

參考資料

- Andrew, S. E., Goldberg Y. P., Kremer, B., Telenius, H., Theimann, J., Adam, S., Starr, E., Squitieri, F., Lin, B., Kalchman, M. A., Graham, R. K., and Hayden, M. R. (1993) The relationship between trinucleotide (CAG) repeat length and clinical features of Huntington's disease. *Nature Genet.* 4, 398-403.
- Aslanidis, C., Jansen, G., Amemiya, C., Shutler, G., Mahadevan, M., Tsilfidis, C. et al. (1992) Cloning of the essential myotonic dystrophy region: mapping of the putative defect. *Nature* 355, 548-551.
- Boucher, C. A., King, S. K., Carey, N., Krahe, R., Winchester, C. L., Rahman, S., Creavin, T., Meghji, P., Bailey M. E. S., Chartier, F. L., Brown, S. D., Siciliano, M. J., and Johnson, K. L. (1995) A novel homeodomain-encoding gene is associated with a large CpG island interrupted by the myotonic dystrophy unstable (CTG)_n repeat. *Hum. Mol. Genet.* 4, 1919-1925.
- Brook, J. D., McCurrach, M. E., Harley, H. G., Buckler, A. J., Church, D., Aburatani, H., Hunter, K., Stanton, V. P., Thirion, J.-P., Hudson, T., Sohn, R., Zemelman, B., Snell, R. G., Rundle, S. A., Crow, S., Davies, J., Shelbourne, P., Buxton, J., Johns, C., Juvonen, V., Johnson, K., Harper, P. S., Shaw, D. J., and Housman D. E. (1992) Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* 68, 799-808.
- Buxton, J., Schelbourne, P., Davies, J., Jones, C., Van Tongeren, T.,

- Aslanidis, C., De Jong, P., Jansen, G., Anvert, M. et al.. (1992)
Detection of an unstable fragment of DNA specific to individuals with
myotonic dystrophy. *Nature* 355, 547-548.
- Davis, B. M., McCurrach, M. E., Taneja, K. L., Singer, R. H., and
Housman, D. E. (1997) Expansion of a CUG trinucleotide repeat in
the 3' untranslated region of myotonic dystrophy protein kinase
transcripts results in nuclear retention of transcripts. *Proc. Natl. Acad.
Sci., USA* 94 (14), 7388-7393.
- Davies, S. W., Turmaine, M., Cozens, B.A., DiFiglia, M., Sharp, A. H.,
Ross, C. A., Scherzinger, E., Wanker, E. E., Mangiarini, L., and Bates,
G. P. (1997) Formation of neuronal intranuclear inclusions underlies
the neurological dysfunction in mice transgenic for the HD mutation.
Cell 90 (3), 537-548.
- Dubowitz, V. (1985) Histological and histochemical stains and reactions.
In Bailliere and Tindall (eds), *Muscle Biology. A Practical Approach*,
London, 29-40.
- Duyao, M., Ambrose, C., Myers, R., Novelletto, A., Persichetti, F.,
Fontali, M., Folstein, S. et al. (1993) Trinucleotide repeat length
instability and age of onset in Huntington's disease. *Nature Genet.* 4,
387-392.
- Faber, P. W., Alter, J. R., MacDonald, M. E., and Hart. A. C. (1999)
Polyglutamine-mediated dysfunction and apoptotic death of a
Caenorhabditis elegans sensory neuron. *Proc. Natl. Sci. Council, USA.*
96 (1), 179-184..
- Fu, Y.-H., Friedman, D. L., Richards, S., Pearlman, J. A., Gibbs, R. A.,
Pizzuti, A., Ashizawa, T., Perryman, M. B., Scariato, G., Fenwick, R.

- G., Jr., and Caskey, C. T. (1993) Decreased expression of myotonin-protein kinase messenger RNA and protein in adult form of myotonic dystrophy. *Science* 260, 235-238.
- Fu, Y.-H., Pizzuti, A., Fenwick, R. G., King, J., Jr., Pajnarayan, S., Dunne, P. W., Dubel, J., Nasser, G. A., Ashizawa, T., de Jong, P., Wieringa, B., Korneluk, R., Perryman, M. B., Epstein, H. F., and Caskey, C. T. (1992) An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science* 255, 1256-1258.
- Gerrit B., Nuria P., Ruben A., Istvan K., Manuel P.-A. and Marek M., (1997) muscleblind, a gene required for photoreceptor differentiation in *Drosophila*, encodes novel nuclear Cys3His-type zinc-finger-containing proteins. *Development* 124, 4321-4331.
- Harley, H. G., Brook, J. D., Rundle, S. A., Crow, S., Reardon, W., Buckler, A. J., Harper, P. S., Housman, D. E., and Shaw, D. J. (1992) Expansion of an unstable DNA region and phenotypic variation in myotonic dystrophy. *Nature* 355, 545-546.
- Harper P. S. (1989) Myotonic dystrophy, 2nd ed., WB Saunders, London.
- Harris, S., Moncrieff, C., and Johnson, K. (1996) Myotonic dystrophy: will the real gene please step forward! *Hum. Mol. Genet.* 5, 1417-1423.
- Heath, S. K., Carne, S., Hoyle, C., Johnson, K. L., and Wells, D. J. (1997) Characterization of expression of mDMAHP, a homeodomain-encoding gene at the murine DM locus. *Hum. Mol. Genet.* 6, 651-657.
- Hiroiyuki, K., Takahisa, T., Kwasi, G. M., Akiko, W., Masato, T., and

- Toshiaki, I. (2000) Profiling of *Caenorhabditis elegans* proteins using two-dimensional gel electrophoresis and matrix assisted laser desorption/ionization-time of flight-mass spectrometry. *Electrophoresis* 21, 1755-1765.
- Hoffman-Radvanyi, H., Lavedan, H., Rabes, J. P., Savoy, D., Duros, C., Johnson, K., and Junien, K. (1993) Myotonic dystrophy: absence of CTG enlarged transcript in congenital forms, and low expression of the normal allele. *Hum. Mol. Genet.* 2, 1263-1266.
- Hsiao, K.-M., Lin, H.-M., Pan, H., Li, T.-C., Chen, S.-S., Jou, S.-B., Chiu, Y.-L., Wu, M.-F., Lin, C.-C., and Li, S.-Y. (1999) Application of FTA sample collection and DNA purification system on the determination of CTG trinucleotide repeat size by PCR-based Southern blotting. *J. Clin. Lab. Anal.* 13, 188-193.
- Hsieh, M., Lin, S. J., Chen, J. F., Lin, H. M., Hsiao, K. M., Li, S. Y., Li, C., and Tsai, C. J. (2000) Identification of the spinocerebellar ataxia type 7 mutation in Taiwan: application of PCR-based Southern blot. *J. Neurol.* 247, 623-629.
- Hsieh-Li, H. M., Chang, J. G., Jong, Y. J., Wu, M. H., Wang, N. M., Tsai, C. H., and Li, H. (2000) A mouse model for spinal muscular atrophy. *Nature Genet.* 24, 66-70.
- Hofferbert, S., Schanen, N. C., Chehab, F., and Francke, U. (1997) Trinucleotide repeats in the human genome: size distributions for all possible triplets and detection of expanded disease alleles in a group of Huntington disease individuals by the Repeat Expansion Detection method. *Hum. Mol. Genet.* 6, 77-83.
- Jansen, G., Groenen, P. J., T. A., Bachner, D., Jap, P. H. K., Coerwinkel,

- M., Oerlemans, F., van den Broek, W., Gohlsch, B., Pette, D., Plomp, J. J., Molenaar, P. C., Nederhoff, M. G. J., van Echteld, C. J. A., Dekker, M., Berns, A., Hameister, H., and Wieronga, B. (1996) Abnormal myotonic dystrophy protein kinase levels produce only mild myopathy in mice. *Nature Genet.* 13, 316-324.
- Jodice, C., Malaspina, P., Persichetti, F., Novelletto, A., Spadaro, M., Giunti, P., Morocutti, C. et al. (1994) Effect of trinucleotide repeat length and parental sex on phenotypic variation in spinocerebellar ataxia 1. *Am. J. Hum. Genet.* 54, 959-965.
- Jou, S.-B., Lin, H.-M., Pan, H., Chiu, Y.-L., Li, S.-Y., and Hsiao, K.-M. (2001). Delineation of CTG repeats and clinical features in Taiwanese Myotonic Dystrophy family. *Proc. Natl. Sci. Council, ROC.* 25, 48-52.
- Kawaguchi, Y., Okamoto, T., Taniwaki, M., Aizawa, M., Inoue, M., Katayama, S., kawakami, H. et al. (1994) CAG expansions in a novel gene from Machado-Joseph Disease at chromosome 14q32.1. *Nature Genet.* 8, 221-227.
- Klesert, T. R., Otten, A. D., Bird, T. D., and Tapscott, S. J. (1997) Trinucleotide repeat expansion at the myotonic dystrophy locus reduces expression of DMAHP. *Nature Genet.* 16, 402-406.
- Klesert, T. R., Cho, D. H., Clark, J. I., Maylie, J., Adelman, J., Snider, L., Yuen, E. C., Soriano, P., and Tapscott, S. J. (2000) Mice deficient in Six5 develop cataracts: implications for myotonic dystrophy. *Nature Genet.* 25, 105-109.
- Koide, R., Ikeuchi, T., Onodera, O., Tanaka, H., Igarashi, S., Endo, K., Takahashi, H., Kondo, R., Ishiawa, A., Hayashi, T., Saito, M., Tomoda,

- A., Miike, T., Naito, H., Ikuta, F., and Tsuji, S. (1994) Unstable expansion of CAG repeat in hereditary dentatorubral-pallidoluysian atrophy (DRPLA). *Nature Genet.* 6, 9-13.
- Koob, M. D., Moseley, M. L., Schut, L. J., Benzow, K. A., Bird, T. D., Day, J. W., and Ranum, L. P. M. (1999) An untranslated CTG expansion causes a novel form of spinocerebellar ataxia. *Nature Genet* 21, 379-384.
- La Spada. A. R., Wilson, E. M., Lubahn, D. B., Harding, A. E., and Fischbeck, K. H. (1991) Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 352, 77-79.
- Lin, M. J. and Lin-Shiau, S. Y. (2000) Suramin protects the murine motor nerves from the toxic effects of presynaptic Ca^{2+} channel inhibitors. *Neurosci. Letters* 287, 97-100.
- Lin-Shiau, S. Y. and Lin, M. J. (1999) Suramin inhibits the toxic effects of presynaptic neurotoxins at the mouse motor nerve terminals. *Euro. J. Pharmacol.* 382, 75-80.
- Lin, M. J. and Lin-Shiau, S. Y. (1999) Enhanced spontaneous transmitter release at murine motor nerve terminals with cyclosporine. *Neuropharmacol.* 38, 195-198.
- Lin, M. J. and Lin-Shiau, S. Y. (1998) Phorbol ester-induced contractions of mouse detrusor muscle are inhibited by nifedipine. *Naunyn-Schmiedeberg's Arch Pharmacol.* 357, 553-557.
- Lin-Shiau, S. Y. and Lin, M. J. (1998) Studies on curare-like action of the tripeptide carbobenzoxy-Gly-Gly-Arg- β -naphthylamide in mouse diaphragm. *Euro. J. Pharmacol.* 343, 51-56 .
- Lin, M. J. and Lin-Shiau, S. Y. (1997) Multiple types of Ca^{2+} channels at

- the mouse motor nerve terminals. *Euro. J. Neurosci.* 9: 817-823 (1997).
- Lin, M. J. and Lin-Shiau, S. Y. (1996) Ruthenium Red, a novel enhancer of K⁺ currents at mouse motor nerve terminals. *Neuropharmacol.* 35, 615-623.
- Lin, M. J. and Lin-Shiau, S. Y. (1995) Functional assessment of Ca²⁺-current in the mouse motor nerve terminals. *Neurosci. Letter* 195, 21-24.
- Lin-Shiau, S. Y., Liu, S. H., and Lin, M.J. (1993) Use of ion channel blockers in the exploration of possible mechanisms involved in the myopathy of diabetic mice. *Naunyn-Schmiedeberg Arch. Pharmacol.* 348: 311-318.
- Liquori, C. L., Ricker, K., Moseley, M., Jacobsen, J. F., Kress, W., Naylor, S., Day, J. W., and Ranum, L. P. W. (2001) Myotonic dystrophy type 2 caused by a CCTG expansion in intron 1 of ZNF9. *Science* 293, 864-867.
- Maciel, P., Gaspar, C., DeStefano, A. L., Silveira, L., Coutinho, P., Radvany, J., Dawson, D. M., Sundarsky, L., Guimaraes, J., Loureiro, J. E. L., Nezarati, M. M., Corwin, L. L., Lopes-Cendes, I., Rooke, K., Rosenberg, R., MacLeod, P., Farrer, L. A., Sequeiros, J., and Rouleau, G. A. (1995) Correlation between CAG repeat length and clinical features in Machado-Joseph disease. *Am. J. Hum. Genet.* 57, 54-61.
- Mahadevan, M., Tsilfidis, C., sabourin, L., Shutler, G., Amemiya, C., Jansen, G., Neville, C., Narang, M., Barcero, J., O'Hoy, K., Leblond, S., Earle-Macdonald, J., de Jong, P. J., Wieringa, B., and Korneluk, R. G. (1992) Myotonic dystrophy mutation: an unstable CTG repeat in

- the 3' untranslated region of the gene. *Science* 255, 1253-1255.
- Mankodi, A., Logigian, E., Callahan, L., McClain, C., White, R., Henderson, D., Krym, M., and Thornton, C., A. (2000) Myotonic dystrophy in transgenic mice expressing an expanded CUG repeat. *Science* 289, 1769-1772.
- Mankodi, A., Urbinati, C. R., Yuan, Q. P., Moxley, R. T., Sansone, V., Krym, M., Henderson, D., Schalling, M., Swanson, M. S., and Thornton, C. A. (2001) Muscleblind localizes to nuclear foci of aberrant RNA in myotonic dystrophy types 1 and 2. *Hum. Mol. Genet.* 10 (19), 2165-2170.
- Martorell, L., Pujana, M. A., Volpini, V., Sanchez, A., Joven, J., Vilella, E., and Estivill, X. (1997) The repeat expansion detection method in the analysis of diseases with CAG/CTG repeat expansion: usefulness and limitations. *Hum. Mut.* 10, 486-488.
- McLaughlin, B. A., Spencer, C., and Eberwine, J. (1996) CAG trinucleotide RNA repeats interact with RNA-binding proteins. *Am. J. Hum. Genet.* 59, 561-569.
- McMurray, C. T. (1999) DNA secondary structure: a common and causative factor for expansion in human disease. *Proc. Natl. Acad. Sci., USA* 96, 1823-1825.
- Michalowski, S., Miller, J. W., Urbinati, C. R., Paliouras, M., Swanson, M. S., and Griffith, J. (1999) Visualization of double-stranded RNAs from the myotonic dystrophy protein kinase gene and interactions with CUG-binding protein. *Nucleic Acids Res.* 27 (17), 3534-3542.
- Miller, J. W., Urbinati, C. R., Teng-umnuay, P., Stenberg, M. G., Bryne, B. J., Thornton, C. A., and Swanson, M. S. (2000) Recruitment of human

- muscleblind proteins to (CUG)_n expressions associated with myotonic dystrophy. *The EMBO J.* 19 (17), 4439-4448.
- Milne, C. A., and Hodgkin, J. (1999) ETR-1, a homologue of a protein linked to myotonic dystrophy, is essential for muscle development in *Caenorhabditis elegans*. *Current Biology* 9, 1243-1246.
- Napierala, M., and Krzyzosiak, W. J. (1997) CUG repeats present in myotonin kinase RNA form metastable 'slippery' hairpins. *J. Biol. Chem.* 272, 31079-31085.
- Nelson, D. L. (1996) Allelic expansion underlies many genetic diseases. *Growth: Genetics & Hormones* 12, 1-4.
- Ordway, J. M., Tallaksen-Greene, S., Gutekunst, C. A., Bernstein, E. M., Cearley, J. A., Wiener, H. W., Dure, L. S., Lindsey, R., Hersch, S. M., Jope, R. S., Albin, R. L., and Detloff, P. J. (1997) Ectopically expressed CAG repeats cause intranuclear inclusions and a progressive late onset neurological phenotype in the mouse. *Cell* 91 (6), 753-763.
- Pan, H. and Griep, A. E. (1994) Altered cell cycle regulation in the lens of HPV-16 E6 or E7 transgenic mice: implications of tumor suppressor gene function in development. *Genes & Dev.* 8, 1285-1299.
- Pan, H. and Griep, A. E. (1995) Temporally distinct patterns of p53-dependent and p53-independent apoptosis during mouse lens development. *Genes & Dev.* 9, 2157-2169.
- Pan, H., Yin, C., and Van Dyke T. A. (1997) Apoptosis and cancer mechanisms. *Cancer Surveys* 29, 305-327.
- Pan, H., Yin, C., Dyson, N.J., Harlow, E., Yamasaki, L., and Van Dyke, T. (1998) Key roles for E2F1 in signaling p53-dependent apoptosis and in

- cell division within developing tumor. *Mol. Cell* 2, 283-292.
- Pan, H., Lin, H.-M., Ku, W.-Y., Li, T.-C., Li, S.-Y., Lin, C. C., and Hsiao, K.-M. (2001) Haplotype analysis of the myotonic dystrophy type 1 (DM1) locus in Taiwan: implications for low prevalence and founder mutations of Taiwanese myotonic dystrophy. *Eu. J. Hum. Genet.* 9, 638-641.
- Pan, H., Li, Y.-Y., Li, T.-C., Tsai, W.-T., Li, S.-Y., and Hsiao, K.-M. Increased (CTG/CAG)_n lengths in the myotonic dystrophy type 1 and Machado-Joseph disease genes in idiopathic azoospermia patients. (revised) *Hum. Repro.*.
- Pan, H., Li, Y.-Y., Tsai, W.-T., Lin, C.-C., and Hsiao, K.-M. Higher frequency of large normal trinucleotide repeat lengths at spinocerebellar ataxia type 8 and Machado-Joseph disease loci in myotonic dystrophy type 1 patients.
- Philips, A.V., Timchenko, L.T., and Cooper, T.A. (1998) Disruption of splicing regulated by a CUG-binding protein in myotonic dystrophy. *Science* 280, 737-741.
- Ruben A., Andreas P., Nuria P., Gerrit B., Ignacio P., Marek M., Manuel P.-A., and Mary K.-B., (1998) The *muscleblind* Gene Participates in the Organization of Z-Bands and Epidermal Attachments of *Drosophila* Muscles and Is Regulated by *Dmef2*. *Developmental Biology* 195, 131–143
- Ranum, L. P. W., Chung, M., Banfi, S., Bryer, A., Schut, L. J., Rameser, R., Duvick, L. A. et al. (1994) Molecular and clinical correlations in spinocerebellar ataxia type 1: evidence for familial effects on the age at onset. *Am. J. Hum. Genet.* 55, 244-252.

- Reddy, S., Smith, D. B. J., Rich, M. M., Leferovich, J. M., Reilly, P., Davis, B. M., Tran, K., rayburn, H., Bronson, R., Cros D., Balice-Gordon, R. J., and Housman, D. (1996) Mice lacking the myotonic dystrophy protein kinase develop a late onset progressive myopathy. *Nature Genet.* 13, 325-335.
- Sarker, P. S., Appukuttan, B., Han, J., Ito, Y., Ai, C., Tsai, W., Chai, Y., Stout, J. T., and Reddy, S. (2000) Heterozygous loss of Six5 in mice is sufficient to cause ocular cataracts. *Nature Genet.* 25, 110-114.
- Satyral, S. H., Schmidt, E., Kitagawa, K., Sondheimer, N., Lindquist, S., Kramer, J. M., and Morimoto, R. I. (2000) Polyglutamine aggregates alter protein folding homeostasis in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci.. USA* 97 (11), 5750-5755.
- Schalling, M., Hudson, T., Buetow, K., and Housman, D. (1993) Direct detection of novel expanded trinucleotide repeats in the human genome. *Nature Genet.* 4,135-139.
- Seznec, H., Agbulut, O., Sergeant, N., Savouret, C., Ghestem, A., Tabti, N., Willer, J. C., Ourth, L., Duros, C., Brisson, E., Fouquet, C., Butler-Browne, G., Delacourte, A., Junien, C., and Gourdon G. (2001) Mice transgenic for the human myotonic dystrophy with expanded CTG repeats display muscular and brain abnormalities. *Hum. Mol. Genet.* 10 (23), 2717-2726.
- Snell, R. G., MacMillan, J. C., Cheadle, J. P., Fenton, I., Lazarou, L. P., Davies, P., MacDonald, M. E., Gusella, J. F., Harper, P. S., and Shaw, D. J. (1993) Relationship between trinucleotide repeat expansion and phenotypic variation in Huntington's disease. *Nature Genet.* 4, 393-397.

- Takano, H., Cancel, G., Ikeuchi, T., Lorenzetti, D., Mawad, R., Stevanin, G., Didierjean, O., Durr, A., Oyake, M., Shimohata, T., Sasaki, R., Koide, R., Igarashi, S., Hayashi, S., Takiyama, Y., Nishizawa, M., Tanaka, H., Zoghbi, H., Brice, A., and Tsuji, S. (1998) Close associations between prevalences of dominantly inherited spinocerebellar ataxias with CAG-repeat expansions and frequencies of large normal CAG alleles in Japanese and Caucasian populations. *Am. J. Hum. Genet.* 63, 1060-1066.
- Taneja, K. L., McCurrach, M., Schalling, M., Housman, D., and Singer, R. H. (1995) Foci of trinucleotide repeat transcripts in nuclei of myotonic dystrophy cells and tissue. *J. Cell Biol.* 128, 995-1002.
- Thornton C. A., Wymer, J. P., Simmons, Z., McClain, C., and Moxley, R. T. (1997) Expansion of the myotonic dystrophy CTG repeat reduces expression of the flanking *DMAHP* gene. *Nature Genet.* 16, 407-409.
- Tian, B., White, R. J., Xia, T., Welle, S., Turner, D. H., Mathews, M. B., and Thornton, C. A. (2000) Expanded CUG repeat RNAs form hairpins that activate the double-stranded RNA-dependent protein kinase PKR. *RNA* 6, 79-87.
- Timchenko, L. T., and Caskey, C. T. (1996) Trinucleotide repeat disorders in humans: discussions of mechanisms and medical issues. *FASEB J.* 10, 1589-1597.
- Timchenko, N.A., Iakova, P., Cai, Z. J., Smith, J. R., and Timchenko, L.T. (2001) Molecular basis for impaired muscle differentiation in myotonic dystrophy. *Mol. Cell. Biol.* 21 (20), 6927-6938.
- Tiscornia, G., and Mahadevan, M. S. (2000) Myotonic dystrophy: the role of the CUG triplet repeats in splicing of a novel DMPK exon and

- altered cytoplasmic DMPK mRNA isoform ratios. *Mol Cell* 5, 959-967.
- Verkerk, A.J.M.H., Pieretti, M., Sutcliffe, J. S., Fu, Y.-H., Kuhl, D. P. A., Pizzuti, A., Reiner, O., Richards, S., Victoria, M. F., Zhang, F., Eussen, B. E., van Ommen, G.-J. B., Blonden, L. A. J., Riggins, G. J., Chastain, J. L., Kunst, C. B., Galjaard, H., Caskey, C. T., Nelson, D. L., Ooster, B. A., and Warren, S. T. (1991) Identification of a gene (*FMR-1*) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 65, 905-914.
- Vincent, J. B., Paterson, A. D., Strong, E., Petronis, A., and Kennedy, J. L. (2000) The unstable trinucleotide repeat story of major psychosis. *Am. J. Med. Genet.* 97, 77-97.
- Wang, Y.-H. and Griffith, J. (1995) Expanded CTG triplet blocks from the myotonic dystrophy gene create the strongest known natural nucleosome positioning elements. *Genomics* 25, 570-573.
- Warrick, J. M., Paulson, H. L., Gray-Board, G. L., Bui, Q. T., Fischbeck, K. H., Pittman, R. N., and Bonini, N. M. (1998) Expanded polyglutamine protein forms nuclear inclusions and causes neural degeneration in *Drosophila*. *Cell* 93 (6), 939-949.
- Wooster, R., Cleton-Jansen, A. M., Collins, N., Mangion, J., Cornelis, R. S., Cooper, C. S., Gusteson, B. A., Ponder, B. A., von Deimling, A., Wiestler, O. D., et al. (1994) Instability of short tandem repeats (microsatellites) in human cancers. *Nature Genet.* 6, 152-156.
- Wu, Y. and Horvitz, H. R. (1998a) *C. elegans* phagocytosis and cell-migration protein CED-5 is similar to human DOCK180. *Nature* 392, 501-504.

Wu, Y. and Horvitz, H. R. (1998b) The *C. elegans* cell corpse engulfment gene *ced-7* encodes a protein similar to ABC transporters. *Cell* 93, 951-960.